




Contents lists available at ScienceDirect

Journal of Volcanology and Geothermal Research

journal homepage: www.elsevier.com/locate/jvolgeores

The significance of maar volcanoes for palaeoclimatic studies in China

Jing Wu^{a,b,*}, Zeyang Zhu^{a,c}, Chunqing Sun^{a,b}, Patrick Rioual^{a,b}, Guoqiang Chu^{a,b}, Jiaqi Liu^{c,a,*}^a Key Laboratory of Cenozoic Geology and Environment, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing 100029, China^b Institutions of Earth Science, Chinese Academy of Sciences, Beijing 100029, China^c University of Chinese Academy of Sciences, Beijing 100049, China

ARTICLE INFO

Article history:

Received 3 April 2018

Received in revised form 26 July 2018

Accepted 7 September 2018

Available online xxxx

Keywords:

Maar lakes

Varves

Palaeoclimate

Tephrochronology

Lacustrine sediments

ABSTRACT

The sediments of maar lakes often provide paleoenvironmental records which are highly suitable for palaeoclimatic studies. The main advantages of the records from maar lakes compared with other lacustrine records are as follows: First, the presence of varves (annually laminated layers) and tephra layers, as well as other materials (like terrestrial plant macrofossils, charcoals, organic carbon, organic shells) suitable for isotopic dating (like AMS ^{14}C , ^{210}Pb , ^{137}Cs , $^{238}\text{U}/^{230}\text{Th}$, $^{40}\text{Ar}/^{39}\text{Ar}$), provides the possibility of developing several independent chronologies, enabling the development of robust, high-resolution chronologies. Second, maars and maar lakes can provide continuous sedimentary records spanning tens of thousands of years (Tianyang Maar in China is hosted the longest known record spanning at least 400 kyr BP) that are ideal for the investigation of climate change on different timescales, including orbital, millennial-centennial and annual-decadal. Third, recent studies of biomarkers based on long-chain alkenones (LCAs) and glycerol dialkyl glycerol tetraethers (GDGTs) have demonstrated the potential of maar lakes for quantitative paleotemperature reconstruction. In this study, we review some of the more significant achievements of palaeoclimatic research based on maar lake sediment sequences from China, with the focus on chronology, multiple timescales and quantitative paleoclimatic reconstruction.

© 2018 Elsevier B.V. All rights reserved.

Contents

1. Introduction	0
2. Study areas	0
3. Chronology	0
3.1. Varve chronology	0
3.2. Radioisotope dating	0
3.3. Tephrochronology	0
4. Climate change on multiple timescales	0
4.1. Modern climate in the context of long-term orbitally-forced climate change	0
4.2. Millennial-scale abrupt climate events and their forcing mechanisms	0
4.3. Prediction of future climate change on annual to decadal scales	0
5. Quantitative paleotemperature reconstructions	0
6. Summary	0
Acknowledgments	0
References	0

1. Introduction

Maar-diatreme volcanoes are small volume monogenetic but commonly complex volcanoes formed in a relatively short time through phreatomagmatic explosive eruptions resulted and excavated and collapsed crater – the maar – cut into the syn-eruptive surface (Lorenz,

* Corresponding authors at: Key Laboratory of Cenozoic Geology and Environment, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing 100029, China.
E-mail addresses: wujing@mail.igcas.ac.cn (J. Wu), liujq@mail.igcas.ac.cn (J. Liu).

1986; White and Ross, 2011; Németh and Kereszturi, 2015). The exact process behind the formation of the maar crater is under focused research in the last decade resulting two distinct endmember of explosive origin; (i) incremental growth of the crater vs. (ii) gradual growth by multi-explosive processes in near surface areas (White and Ross, 2011; Valentine et al., 2017). After the formation of the maar crater, it become the ideal terrestrial closed sedimentary basin (Negendank and Zolitschka, 1993a) that has a small surface area and in comparison to that deep basin where following the tephra ring erosion, quite long-lasting lacustrine sedimentation can take place (Pirrung et al., 2003).

Maar-diatreme structures can host climate-sensitive lakes in which long and continuous sedimentary sequences can accumulate. This is one of the reasons for the rapid increase of studies on maar-diatreme volcanoes (Creer, 1996; Creer and Morris, 1996; Marchetto et al., 2015; Turner, 1998; Sirocko, 2016; Zolitschka et al., 2000, 2013). Maar lakes are now widely acknowledged to be of great significance in the research field of palaeoclimatic change research on the continents. In the past decade intensive paleoclimatic researches were conducted also in the southern hemisphere including maars in Argentina (e.g. Zolitschka et al., 2013) and New Zealand (e.g. Augustinus, 2007).

Palaeoclimatic research is a major topic in earth science. In recent decades, studies of deep-sea sediments, and ice and loess deposits have demonstrated the occurrence of orbitally-forced climatic cycles on multi-millennial timescales, which have confirmed that orbital forcing is a major driver of large-scale changes in earth's climate and environment (Berger, 1988). Later research revealed that the climate system also exhibits higher-frequency changes and abrupt events on various timescales, ranging from millennial to annual (Bond et al., 1997). A major objective of global change research is to forecast future trends in climate, especially over the next few decades, and to achieve these goals, palaeoclimatic records should be of interdecadal to annual resolution. The longest global observed temperature records only span the last ~150 years (Brohan et al., 2006), and therefore they are of limited usefulness for assessing climate changes on interdecadal or longer scales. This limitation hinders our understanding of the mechanisms behind both gradual and abrupt climate change, and consequently there is an ongoing search for continuous, long-duration, high resolution geological archives for which multi-proxy environmental indicators are available (Creer, 1996; Creer and Morris, 1996; Garcin et al., 2006; Augustinus et al., 2012; Zaarur et al., 2018). In this context, the sedimentary record of maar lakes, as a continental archive, make a significant contribution.

Maar lakes are formed by phreatomagmatic volcanic eruptions; they consist of a crater cut into the pre-eruptive ground, surrounded by an ejecta ring, underlain by a diatreme (Lorenz, 1973, 2003, 2007; Negendank and Zolitschka, 1993b; Liu et al., 1996; Valentine, 2012). Maar-diatreme structures can host climatically-sensitive lakes which preserve long and continuous sedimentary sequences, which is one of the main reasons for the rapid increase in the number of studies of maar-diatreme volcanoes (White and Ross, 2011; Zolitschka et al., 2000), and maar lakes are now widely acknowledged to be of high significance for palaeoclimatic research on continents. The major advantages of maar lakes for paleoenvironmental studies can be summarized as follows:

- (i) Maar lakes generally do not have inflows and outflows and the water level is mainly controlled by groundwater influx (Büchel, 1993); thus, the hydrology and sedimentation processes of maar lakes are relatively simple to interpret in terms of climatic change, compared with lakes with more complex hydrological settings.
- (ii) The size, original depth and shape of maar craters result in a relatively low surface area/depth ratio, which limits the effects of waves on the sedimentary record, providing suitable conditions for the preservation of annual laminations or varves (Graetinger, 2018).

- (iii) The bottom waters of some deep maar lakes are hypoxic, which substantially limits the number and activity of benthic organisms; the resulting suppression of bioturbation also promotes the preservation of the protogenic structure of the sediment (Zolitschka et al., 2015).
- (iv) The sediments of maar lakes provide a highly diverse and complementary suite of proxy climatic indicators. They include biological indicators such as pollen derived from the terrestrial vegetation, and numerous aquatic organisms such as diatoms, chrysophytes, chironomids, ostracods; and physical and chemical parameters such as magnetic properties, and the stable isotopes of oxygen and carbon (Marchetto et al., 2015). Cross-comparison of different indicators analyzed from the same sedimentary sequence enables robust and reliable multi-proxy-based palaeoclimate reconstructions to be produced.

In summary, maar lakes provide optimal conditions for the accumulation and preservation of high-resolution sedimentary records on continents, and these natural archives are of high value for multi-proxy studies of palaeoenvironmental and palaeoclimatic changes.

2. Study areas

Prior to the 1980s, Europe was the main study region for maars and pioneering research was carried out in the West-Eifel Volcanic Field, in northwest Germany (Ollier, 1967; Lorenz, 1975; Büchel and Lorenz, 1982; Negendank, 1984). At that time, research focused on the structure, formation and geology of maars rather than on their climatic records. In the late 1980s, the International Geosphere-Biosphere Programme (IGBP) advocated the search for more high-resolution terrestrial sedimentary records worldwide, and to this end various research agencies in Europe launched and funded a series of drilling programs, such as GEO-MAARS, EURO-MAARS, the European Lake Drilling Program (ELDP) and the ELSA Project (Eifel Laminated Sediment Archive) (Büchel, 1993; Creer, 1996; Sirocko et al., 2013; Sirocko, 2016). Consequently, research on maar lakes moved from Germany to Italy, and records from maar lakes in central and southern Italy provide continuous lacustrine sedimentary records back to at least 70 kyr BP (Creer and Morris, 1996). At the end of the 20th century, there was a global expansion of maar lake research, which extended the regions of study to central France and central Spain, and sporadic studies were also reported from Asia, Africa and Australia (Augustinus, 2012; Barker et al., 2003; D'Costa and Kershaw, 1995; Juvigné et al., 1993; Liu et al., 1996; Ortiz et al., 2013). In addition, the Laguna Potrok Aike Scientific Drilling Project (PASADO) was conducted in South America, in the southernmost tip of Argentina (Zolitschka et al., 2009). Locations and details of research projects on maar lakes are provided in Fig. 1 and Table 1. Numerous continuous sedimentary records from maar lakes have been obtained, with important findings in the fields of chronology (e.g. Brauer et al., 1999; Lane et al., 2013), sedimentology (e.g. Giguet-Covex et al., 2010; Chu et al., 2013), geochemistry (e.g. Parplies et al., 2008), micropaleontology (e.g. Kattel and Sirocko, 2011), and paleomagnetism (e.g. Creer and Morris, 1996) as the most common research focuses. Thus, maar lakes are exhibiting great potential for addressing significant questions about long-term terrestrial climatic change.

In China, research on maar lakes began in 1996 (Liu et al., 1996). The main study areas are the Leiqiong Volcanic Field in South China and the Longgang Volcanic Field in Northeast China (Fig. 2). The Leiqiong Volcanic Field is part of the epi-continental rift volcanic belt that crosses Leizhou Strait in Southern China (Fig. 2a). The volcanic field is characterized by a variety of basaltic volcanoes, typical lava flow structures and long lava tunnels (the longest one is longer than 12 km) (Tao, 2007). Volcanic activity began in the Oligocene and lasted through the Holocene. Early volcanism was dominated by quartz tholeiites and olivine




Fig. 1. Location of the maars and maar lakes with reported palaeoclimatic records. (Modified from Liu et al., 2013).

tholeiites, and late volcanism was dominated by alkali olivine basalts and basanites (Huang et al., 2007). There are numerous maars located in this volcanic field: for example, Huguangyan (21°09'N, 110°17'E), Tianyang (20°31'N, 110°18'E), Jiudouyang (20°38'N, 110°02'E), Qingtongyang (20°34'N, 110°10'E), and Shuangchiling Maar (19°57'N, 110°11'E) (Hunag et al., 1993; Mingram et al., 2000; Fuhrmann et al., 2003; Yang et al., 2009). Most are dry maars, except for Huguangyan and Shuangchiling Maar Lake. Tianyang Maar is one of the largest craters in China, with a diameter of ~3000 m and a maximum depth of 223 m, and was formed by at least 400 kyr BP (Zheng and Guiot, 1999).

Longgang Volcanic Field located in southeastern Jilin province (Northeast China) has a length of 85 km from south to north, a width of 50 km from east to west, an area of about 4250 km², and an area covered by volcanic rocks of approximately 1700 km² (Fan et al., 1999, 2002). The basement is mainly composed of Pre-Cambrian basement rocks, such as Archean migmatitic gneiss, granite and leptynite in the Anshan group. The distribution of volcanoes generally follows alignments associated with major structural lineaments of the basement (NWW or near EW), always at the intersection of NE and EW, part of the East Asian continental rift system (Liu et al., 2009). Quaternary volcanic activity in Longgang Volcanic Field produced basaltic magmatic and phreatomagmatic eruptions, which formed at least 160 monogenetic volcanoes. Eight maar lakes (Sihailongwan, Donglongwan, Xiaolongwan, Sanjiaolongwan, Erlongwan, Donglongwan, Nanlongwan and Longquanlongwan) are located in this volcanic field (Fig. 2b). It is a key region for research on regional climate change controlled by the East Asian Monsoon. The tephra rings of these maars typically consists of pyroclastic beds of base surges, lava flows and scoria fall deposits (Fan et al., 1999, 2002).

After >20 years' research, studies of Chinese maars have progressed substantially, especially in the areas of geochronology, palaeoclimatology, and currently their volcanology. We now review progress in first two major research areas.

3. Chronology

A reliable chronology is a fundamental requirement for palaeoclimate studies and is the basis for local, regional and global correlation of climatic events and for determining phase relationships between signals. The presence of laminations in maar sediments

provides the most robust chronological framework, providing the annual character of the laminations is confirmed (Zolitschka et al., 2015). If not, a reliable chronological framework can still be established based on isotope dating or tephrochronology.

3.1. Varve chronology

The term 'varve' is used to describe all types of annually laminated layers deposited in terrestrial and marine settings (De Geer, 1912; Zolitschka et al., 2015). The most significant aspect of a varve chronology for palaeoclimatic and paleoenvironmental research is that it provides precise dating for different temporal scales and for specific events, which can then be used as isochrones on a regional or global scale (Nakagawa et al., 2012; Zolitschka et al., 2015). Varves were first recognized and defined in glacial lake sediments from high latitude regions (De Geer, 1912), where there are pronounced seasonal variations in climatic and depositional processes which result in stronger contrasts than is the case for mid- and low latitude lakes. Despite the disadvantage of less pronounced seasonal changes in climate, maar lakes located in mid- and low latitude areas still provide favorable conditions (low surface area/depth ratio, deep water with a stratified water column and hypoxic bottom water) which facilitate the formation and preservation of varves. Thus, maar lake sediments with varves are considered among the most valuable environmental archives of past climatic changes in continental areas.

Different types of varve are recognized in maar lake sediments according to their composition, and there are three main types: clastic, biogenic and endogenic (Table 2). Among them, biogenic or clastic-biogenic varves have the widest distribution. Well-known examples of this type of varve occur in Lake Barrine in Australia (Walker, 2011; Walker and Owen, 1999), Lac Pavin and Lac du Bouchet in France (Giguet-Covex et al., 2010; Stebich et al., 2005), Lake Albano in Italy (Lami et al., 1994), and Holzmaar (Zolitschka, 1998) and Meerfelder Maar (Brauer et al., 1999; Lane et al., 2015; Martin-Puertas et al., 2017) in Germany. Calcareous (endogenic) varves occur in Lake Twintaung in Myanmar (Sun et al., 2016b), and Lake Hoya La Alberca and Lake Hoya Rincón de Paranguico in Mexico (Kienel et al., 2009). Until now, the longest continuous absolute varve chronology established from a maar lake is that for Holzmaar in the West-Eifel Volcanic Field in Germany, which spans ~23,220 years (Zolitschka, 1998).

Table 1

Details of the maars and maar lakes referenced in the text.

Site	Latitude	Longitude	Volcanic field	Country	Interval of study (year BP)	Reference
Lake Holzmaar	50°07'N	6°53'E	West Eifel Volcanic Field	Germany	0–23,220	Zolitschka, 1998; Zolitschka et al., 2000
Lake Meerfelder Maar	50°06'N	6°45'E			0–12,700	Brauer et al., 1999
Lake Schalkenmehrener Maar	50°10'N	6°51'E			0–16,500	Kattel and Sirocko, 2011
Lake Ulmener Maar	50°13'N	6°59'E	Chaîne des Puys	France	0–10,895	Zolitschka et al., 1995; Sirocko et al., 2013
Eckfeld Maar	50°07'N	6°49'E			Eocene	Mingram, 1998
Alleret Maar	45°11'N	3°28'E			550,000–725,000	Nomade et al., 2010
Lac Pavin	45°30'N	2°53'E	Colli Albani Complex	Italy	0–700	Stebich et al., 2005
Lake Albano	41°45'N	12°40'E			0–11,480	Ariztegui et al., 2001
Lagaccione Maar	42°34'N	11°51'E			0–100,000	Magri, 1999
Lago di Mezzano	42°37'N	11°46'E	Monte Vulture		0–34,000	Ramrath et al., 1999, 2000
Lago Grande di Monticchio	40°56'N	15°36'E			0–75,000	Ramrath et al., 1999, 2000; Creer and Morris, 1996; Wulf et al., 2004
Lago di Vico	42°19'N	12°10'E	Vico-Cimino Complex	Hungary	0–90,000	Frank, 1969; Narcisi, 2001
Pula Maar	46°59'N	17°36'E	Bakony-Balaton Highland Volcanic Field		2,600,000–3000,000	Willis et al., 1999
Fuentillejo Maar	38°56'N	4°03'W	Calatrava Volcanic Field	Spain	0–220,000	Ortiz et al., 2013
Lake Tritrivakely	19°47'S	46°55'E	Ankaratra Volcanic Field	Madagascar	0–46,000	Williamson et al., 1998; Disnar et al., 2005
Lake Masoko	9°20'S	33°45'E	Rungwe Volcanic Field	Tanzania	0–50,000	Gibert et al., 2002; Barker et al., 2003; Garcin et al., 2006
Lake Tilo	7°04'N	38°06'E	Bilate River Field	Ethiopian	0–9000	Telford and Lamb, 1999; Lamb et al., 2000
Lake Barombi Mbo	4°40'N	9°24'E	West Cameroon Volcanic Field	Cameroon	0–25,000	Giresse et al., 1994
Laguna Potrok Aike	51°58'S	70°23'W	Pali-Aike Volcanic Field	Argentina	0–51,000	Kliem et al., 2013; Zolitschka et al., 2013
Lago Verde	18°37'N	95°21'W	Tuxtla Volcanic Field	Mexico	0–150	Ruiz-Fernández et al., 2007
Hoya San Nicolás	20°23'N	101°15'W	Michoacán-Guanajuato Volcanic Field		0–11,600	Chaparro et al., 2008; Park et al., 2010
Hoya Rincón de Parangueo	20°26'N	101°15'W			0–9600	Kienel et al., 2009; Park et al., 2010;
Hoya La Alberca	20°23'N	101°12'W			–23–98	Kienel et al., 2009
Zagoskin Lake	63°27'N	162°6'W	St. Michael Volcanic Field	America	0–41,000	Muhs et al., 2003
Lake Terang	38°15'S	142°55'E	Newer Volcanics	Australia	25,000–75,000	D'Costa and Kershaw, 1995
Lake Keilambete	38°12'S	142°53'E	Province		–40–110	Jones et al., 2001; Wilkins et al., 2012
Lake Gnotuk	38°13'S	143°06'E			–40–110	Jones et al., 2001; Wilkins et al., 2012
Lake Bullenmerri	38°15'S	143°06'E			–40–110	Jones et al., 2001
Lake Barrine	17°15'S	145°38'E	Atherton Volcanic Province		0–9000	Dimitriadis and Cranston, 2001; Walker, 2007
Foulden Maar	45°32'S	170°13'E	Waipiata Volcanic Field	New Zealand	Oligocene-Miocene transition	Lindqvist and Lee, 2009
Onepoto Maar	36°48'S	174°45'E	Auckland Volcanic Field		7000–30,000	Augustinus et al., 2012; Sikes et al., 2013
Lake Pupuke	36°47'S	174°46'E			24,500–54,000	Nilsson et al., 2011
Lake Ichi-no-Megata	39°57'N	139°44'E	Megata Volcanic Group	Japan	0–31,000	Okuno et al., 2011
Lake Ni-no-Megata	39°57'N	139°43'E			0–2900	Yamada et al., 2010
Lake San-no-Megata	39°56'N	139°42'E			0–2250	Yamada et al., 2010
Lake Twintaung	22°22'N	95°02'E	Lower Chindwin Volcanic Field	Myanmar	0–530	Sun et al., 2016a, 2016b
Tianyang Maar	20°31'N	110°18'E	Leiqiong Volcanic Field	China	0–400,000	Zheng and Lei, 1999; Zheng and Guiot, 1999
Lake Hugangyan	21°09'N	110°17'E			0–78,000	Liu et al., 2000; Chu et al., 2002; Chu et al., 2017; Mingram et al., 2004b; Liu et al., 2005a; Wang et al., 2007, 2012; Yancheva et al., 2007; Wang et al., 2008, 2012; Hu et al., 2015, 2016
Lake Shuangchiling	19°57'N	110°11'E	Longgang Volcanic Field		0–9000	Yang et al., 2009
Lake Sihailongwan	42°17'N	126°36'E			0–16,700	Chu et al., 2000, 2005a, 2005b, 2011; Liu et al., 2005b; Schettler et al., 2006a, 2006b, 2006c; Parplies et al., 2008; Stebich et al., 2009, 2015
Lake Xiaolongwan	42°18'N	126°21'E			0–1600	Chu et al., 2008, 2009, 2013
Lake Erlongwan	42°18'N	126°21'E			0–13,000	You et al., 2008; You and Liu, 2012

Because of its high quality, the Holzmaar chronology is regarded as a standard stratigraphic sequence for the Late-glacial in Europe, which can be used to supplement or replace the chronologies derived from Greenland ice cores (Litt et al., 2001; Litt and Stebich, 1999). This illustrates the great potential of maar lakes in Europe in establishing highly robust varve chronologies.

In China, absolute varve chronologies have been established from maar lakes in Longgang Volcanic Field, Northeast China, such as Lake Sihailongwan, Erlongwan and Xiaolongwan, and their duration exceeds 14,000 years (Chu et al., 2009; Schettler et al., 2006b; You et al., 2008). In these three lakes, the varves are of the clastic-biogenic type. Based on the dominant types of algae contained in the sediments, the biogenic varves of Lake Sihailongwan can be classified into Chrysophyte and diatom-biogenic varves (Chu et al., 2005a); those of Lake Erlongwan

into Dinocyst and diatom-biogenic varves (You et al., 2008); and those of Lake Xiaolongwan into Dinocyst and Chrysophyte-biogenic varves (Chu et al., 2009). Until now, they represent the only absolute varve chronologies in China that are older than 10 kyr BP.

3.2. Radioisotope dating

Radioisotopic dating methods are necessary for the establishment of a high-resolution chronology, even in maar lakes with varved sediments. Radiocarbon dating is the most widely used dating method for establishing chronologies for maar lake sediments. The bottom waters of maar lakes are typically depleted in dissolved oxygen, which favors the preservation of terrestrial plant remains in the sediments. The use of these terrestrial plant remains potentially avoids the 'hard water'




Fig. 2. Locations of maar volcanoes in China mentioned in text. a. Leiqiong Volcanic Field; b. Longgang Volcanic Field.

and 'carbon reservoir' effects and maximizes the accuracy of radiocarbon dating (Sirocko et al., 2013). The use of accelerator mass spectrometry (AMS) radiocarbon dating considerably reduces the required

minimum mass of the samples to be dated and allows the much wider application of the technique. Radiocarbon dating may be used as the primary method for establish chronologies, as at Fuentillejo maar lake in

Table 2

Details of the varve and other chronologies available for the maars and maar lakes referenced in the text.

Name of maar	Country	Type of chronology	Chronology	Varve type	Varve years	Reference
Holzmaar	Germany	Absolute	Varve, ^{14}C , optically stimulated luminescence, thermoluminescence, paleomagnetism, tephra	Various	23,220	Zolitschka, 1998;
Meerfelder Maar	Germany	Floating	Varve, ^{14}C , paleomagnetism, tephra	Clastic-biogenic	12,700	Zolitschka et al., 2000
Lac Pavin	France	Absolute	Varve	Biogenic (diatom)	700	Brauer et al., 1999; 2000
Lake Albano	Italy	Absolute	Varve/ ^{14}C , tephra, pollen stratigraphy	Biogenic (calcite)	46	Stebich et al., 2005
Lake Hoya La Alberca	Mexico	Floating	Varve, ^{210}Pb , tephra	Calcareous	121	Lami et al., 1994;
Lake Hoya Rincón de Parangueo	Mexico	Floating	Varve, ^{14}C , ^{210}Pb , tephra	Calcareous	104	Ariztegui et al., 2001
Lake Barrine	Australia	Absolute	^{14}C , ^{210}Pb	Detritus poor and - rich calcareous	3000	Kienel et al., 2009
Lake Twintaung	Myanmar	Absolute	Varve, ^{210}Pb , ^{137}Cs , ^{14}C	Clastic-biogenic (dinoflagellate) cysts	530	Kienel et al., 2009
Lake Xiaolongwan	China	Absolute	Varve, ^{14}C , ^{210}Pb , ^{137}Cs , tephra	Biogenic (clastic)	1500	Walker, 2011
Lake Erlongwan	China	Absolute	Varve, ^{14}C , ^{137}Cs	Biogenic (clastic)	12,766	Sun et al., 2016b
Lake Sihailongwan	China	Absolute	Varve, ^{14}C , tephra	Clastic-biogenic	13,759	Chu et al., 2009
						You et al., 2008
						Schettler et al., 2006b

Central Spain (Ortiz et al., 2013), Zagoskin maar lake in western Alaska (Muhs et al., 2003) and Maar Lake Massoko in Tanzania (Gibert et al., 2002), but it is also used to verify varve chronologies from maar lakes, as in Holzmaar in Germany (Hajdas et al., 1995) and Lac du Bouchet in France (Williams et al., 1998). Due to the limitations of the radioactive half-life of ^{14}C , the maximum limit of radiocarbon dating is ~50 kyr BP, and to establish a chronology for older sediment sequences other dating methods are needed.

Other radioisotopic dating methods, such as ^{210}Pb and ^{137}Cs , are generally used for palaeoclimatic reconstruction on short timescales, as in the case of Hoya La Alberca and Hoya Rincón de Parangueo maar lakes in Mexico (Kienel et al., 2009), or for dating the uppermost part of the sedimentary sequence (Sirocko et al., 2013).

In China, radiocarbon dating has been the primary method used for building a chronology at Huguangyan Maar Lake in the Leiqiong Volcanic Field (Mingram et al., 2004b), and was also used to verify the varve chronologies established for Sihailongwan and Erlongwan Maar Lakes in Longgang Volcanic Field (Schettler et al., 2006b; You et al., 2008). In addition, ^{210}Pb and ^{137}Cs dating were used to date the upper part of the sedimentary sequences of several maar lakes in Longgang Volcanic Field (Chu et al., 2005a, 2008; You et al., 2008), and for comparison with the results derived from sediment trap sampling with the aim of determining whether the laminations were annual.

3.3. Tephrochronology

Tephra includes all unconsolidated pyroclasts sourced from explosive volcanic eruptions in various grain sizes: ash (grains < 2 mm in diameter), lapilli (2–64 mm), and blocks (>64 mm) (White and Houghton, 2006). The primary tephra deposits from an eruption essentially have the same near-instantaneous age everywhere they occur, which form isochrons very soon after the eruption (Lowe, 2011). Therefore, the primary definition of tephrochronology is the formation of isochronous marker beds, which are used to connect or correlate sedimentary sequences (marine and lacustrine sediments, ice deposits and other continental sequences) within the area of distribution of the tephra (Lowe, 2011). Tephra layers can be well preserved in maar sediments. If the ages of the tephra layers are known from modern observation, historical documents, ^{40}Ar – ^{39}Ar dating or other dating methods, they can provide absolute dating points for the chronology (Liu et al., 2018; Lowe, 2011). Alternatively, the ages of some tephra layers can also be deduced based on the pre-existing chronology of the corresponding sedimentary archive. In addition, because of the isochronous character of tephra layers, they are very useful for synchronizing regional climatic and environmental changes, once they are recognized and characterized, in various regional sedimentary archives (Lane et al., 2013).

The results of tephrochronology on water-filled and dry maars in the west Eifel volcanic field in Germany show that the tephra layers were produced by several large eruptions during the Middle and Late Pleistocene (Ulmener Maar – 10,904 varve years BP; Laacher See – 12,880 varve years BP; Neapolitan Yellow Tuff/Campi Flegrei volcano – 14 kyr BP; Dümpel Maar – 116 kyr BP; Glees Maar – 151 kyr BP) and they can be used as isochrons for inter-core and inter-site correlations (Brauer et al., 1999; Lane et al., 2015; Sirocko et al., 2013). Among them, the Ulmener Maar Tephra (UMT) and Laacher See Tephra (LST) have been detected in sediment cores from Meerfelder Maar, Schalkenmehrener Maar, Holzmaar, Dehner Maar and Lago Grande di Monticchio, and are important correlation markers between Holocene and Late-glacial core segments from maar lakes in Europe (Lane et al., 2012, 2015; Wulf et al., 2004). Other tephra layers provide additional chronological reference points and the isochrones are either close to or exceed the age limit of ^{14}C dating in Europe and South America. Examples are the Campanian Ignimbrite (CI) Tephra (~39 kyr BP) detected in Lago Grande di Monticchio in Italy (Wulf et al., 2004); the tephra layers detected in Lac du Bouchet Maar and Alleret Maar in France,

which were emitted by the Mont-Dore/Sancy strato-volcano (Pastre et al., 2007); and the lowermost tephra layers detected in Laguna Potrok Aike in Argentina, derived from eruptions of Mt. Burney (Wastegård et al., 2013).

In East Asia, there are several widespread tephra layers that can be traced from Japan to northeast China, such as the Aira-Tn tephra (AT, ~30 kyr BP), the ash from the Changbaishan Millennium eruption (also called the Baitoushan-Tomakomai tephra, B-Tm, AD 946), and the ash from the Changbaishan Qixiangzhan eruption (QEA, ~8100 cal yr BP) (Machida and Arai, 1983; Mingram et al., 2008; Sun et al., 2015, 2016a, 2018). The AT and B-Tm tephra layers can be correlated from maar lake Ichi-no-Megata in Japan to Sihailongwan in China; in addition, the B-Tm tephra is present in Greenland ice cores (Mingram et al., 2008; Okuno et al., 2011; Sun et al., 2014, 2015, 2016a). Thus, these tephra layers are regional marker horizons for palaeoclimatic studies. Sihailongwan Maar Lake also preserves several local basaltic tephra layers, and the varved sediments can be used to determine the relative timing of these eruptions (Liu et al., 2009; Mingram et al., 2004a).

In summary, the characteristics of the deposits that compose maar sediments enable the application of multiple dating methods, which greatly enhances the precision of the chronologies of the sedimentary sequences from maar lakes and dry maars.

4. Climate change on multiple timescales

The maar sedimentary archives enable the decipherment of natural climatic variability from annual to orbital scale (Fig. 3), especially in Quaternary. They also have great potential for providing precise and accurate inter-archive correlations (Vos et al., 2000), for recording cyclical and high-frequency climate signals (Lenz et al., 2010, 2017; Willis and Braun, 1997; Willis et al., 1999), and for identifying the physical mechanisms responsible for the observed changes in climate (Marchetto et al., 2015). Besides Quaternary sites, some older maars provide sedimentary records for palaeoclimatic researches from Pliocene to Eocene (Fox et al., 2010, 2011; Mezger et al., 2013; Németh et al., 2008; Richter et al., 2017; Sabol et al., 2004), based on macrofossils such as plant fossils (Bannister et al., 2012; Conran et al., 2014; Grein et al., 2011; Reichgelt et al., 2013) and animal fossils (De Soler et al., 2012; Kaulfuss et al., 2011; Vass et al., 2000; Wappler et al., 2011), and some microfossils like pollen, algae and coprolites (Conran et al., 2014; Goth and Suhr, 2003; Lenz et al., 2011, 2014).

4.1. Modern climate in the context of long-term orbitally-forced climate change

According to the Milankovitch theory (Milankovitch, 1941; Berger, 1988), glacial-interglacial cycles are primarily forced by summer insolation variations at high latitudes of the Northern Hemisphere. Records from deep-sea sediment cores, ice cores, loess-paleosol profiles and stalagmites, reveal that the cyclical climate changes ranging from ~20 to ~100 kyr in length are caused by variations in Earth orbital eccentricity, obliquity and precession (Fig. 3a). Spectral analysis of a biomarker record from Fuentillejo maar in Spain, spanning the last 220 kyr, revealed significant periodicities at 103, 41, 23 and 19 kyr, which provides strong evidence for the validity of the Milankovitch theory (Ortiz et al., 2013). However, the mechanisms behind the development of glacial stages and their timing remain controversial, and are one of the most challenging issues in palaeoclimate research. Besides summer insolation variations at high latitudes of the Northern Hemisphere, changes in the sea surface temperature (SST) of the tropical oceans and in the carbon reservoir are also regarded as forcing factors of glacial-interglacial cycles (Lea et al., 2000; Wang et al., 2003).

Based solely on Milankovitch theory, the duration of interglacials should be about 10 kyr, which means that the present interglacial, which has already lasted for 11.5 kyr, should have ended about 1.5 kyr ago, and a new glacial stage should have been initiated (e.g. Broecker,




Fig. 3. Paleoclimatic records from maar lakes compared with the marine $\delta^{18}\text{O}$ record. a) Pacific benthic $\delta^{18}\text{O}$ record (Lisiecki and Raymo, 2009). b) Non-arboreal pollen (NAP) record from Tianyang Maar, plotted on an orbital time scale (Zheng and Lei, 1999). c) Total organic matter (TOC) flux from Hugangyan Maar Lake (Mingram et al., 2004a, 2004b). d) Record of titanium content from Hugangyan Maar Lake. e) Centennial-millennial-scale record of titanium content from Hugangyan Maar Lake (Yancheva et al., 2007). Dark shading indicates glacial periods and cold events such as the Younger Dryas (YD).

1966). However, others have suggested that the duration of the present interglacial will be 40–50 kyr, or even 100 kyr, because of the elevated levels of atmospheric CO_2 and its effects on Arctic ice sheets (Berger and Loutre, 2002; Ganopolski et al., 2016; Hao et al., 2012). Investigations of maar lake sedimentary sequences in Italy (Lago di Vico) and France (Lac du Bouchet, Praclaux, Ribains) showed that they recorded climatic variations from the present to at least the penultimate glacial period (the Saalian glacial) (Follieri et al., 1998; Turner, 1998), and these records may help to address questions regarding the length of glacial/interglacial cycles. These long records can provide further information on the last interglacial period, which is regarded as an analogue of the present interglacial and may help us to properly understand the position of modern climate in glacial/interglacial cycles. A multi-proxy lacustrine record of climate change during the last interglacial period, based on diatom and pollen analysis, from Ribains maar in France, indicates that the climate in central and southern Europe was warm and stable during the first 10–12 kyr of the last interglacial period, and somewhat colder but still stable during the subsequent ~10 kyr (Rioual et al., 2007). This implies that the climate of the current interglacial may become somewhat colder but will still remain stable and last for a further 10 kyr (Rioual et al., 2007).

In China, the continuous sedimentary record from Tianyang Maar (Leiqiong Volcanic Field/South China) spans at least the last 400 kyr (Zheng and Guiot, 1999, Fig. 3b). A quantitative pollen-based temperature record from the site provides an improved correlation between terrestrial and marine records, such as the records of sea-level change and SST in the adjacent South China Sea (Zheng and Lei, 1999). This correlation suggests that the low-latitude tropical climate in China responded sensitively to global ice volume and/or sea level changes; however, higher resolution studies of the long sedimentary records from maars

in China are needed. Undoubtedly, such studies will improve our understanding of glacial-interglacial cycles and the modern climate in the context of orbital forcing.

4.2. Millennial-scale abrupt climate events and their forcing mechanisms

Climatic instability is characteristic of the Late Pleistocene and the Holocene. During the Late Pleistocene, climatic variability on a millennial scale is evidenced by Dansgaard-Oeschger (D-O) cycles (Dansgaard et al., 1993), which are detected in marine sediments and ice cores (Bond et al., 1997). The Holocene equivalent of D-O cycles is a series of climatic shifts with a cyclicity close to 1470 ± 500 years (Bond et al., 1997). The evidence for D-O cycles is the presence of ice-rafted, carbonate-rich debris layers in marine sediments which are interpreted as indicating the extension of ice sheets onto shelf areas (Baumann et al., 1995; Bond and Lotti, 1995). Packets of D-O cycles. Packets of D-O cycles are terminated by a pronounced cold event which in sediment cores from the North Atlantic are represented by ice-rafted detrital layers, termed Heinrich events, which result from massive discharges of icebergs from the Laurentide Ice Sheet (Bond et al., 1992). The corresponding temperature changes in North Atlantic surface waters are amplified by the thermohaline circulation and may impact the atmospheric circulation that has a substantial influence on large areas of the Northern Hemisphere, and even globally. However, the causal mechanism of these events remains disputed. A solar forcing mechanism is supported by the close correlation between inferred changes in the production rates of cosmogenic nuclides ^{14}C and ^{10}Be and by centennial-to-millennial scale changes in drift ice proxies measured in deep-sea sediment cores (Bond et al., 2001). However, the millennial-scale cyclicity of solar activity is most significant at periods

of 1000 and 2115 years, but not at periods of 1400 and 1500 years, which are the most pronounced cycles observed in marine sediment records (Damon and Jirikowic, 1992).

Late Pleistocene glacial events detected in the lacustrine data from three European maar lakes, Lago di Mezzano, Lago Grande di Monticchio in Italy and Fuentillejo in Spain, are potentially correlative with the Heinrich events recorded in North Atlantic sediments (Ortiz et al., 2013; Ramrath et al., 1999). This relationship is interpreted as evidence for the Late Pleistocene climate in Europe being influenced by North Atlantic climatic forcing, which may be related to Dansgaard-Oeschger cycles (Ortiz et al., 2013; Ramrath et al., 1999). In contrast, in the sedimentary record from maar Lake Masoko in Tanzania, in tropical southern Africa (Garcin et al., 2006), there is no evidence of the ~6000-year cycles and ~1500-year cycles which characterize North Atlantic climate and iceberg surges during glacial and post-glacial times (Dansgaard et al., 1993; Grootes et al., 1993; Schulz et al., 1999).

In China, the existence of a common millennial-scale periodicity during the Holocene has been detected in the records of maar lakes in both low and mid-latitude region (Chu et al., 2014; Wang et al., 2016). A ~1500-year cycle detected in the records from maar lakes Huguangyan (1140 years) and Xiaolongwan (1020–1050 and 1750–2041 years) implies that the millennial monsoon cycle in East China is associated with sea surface temperature (SST) variations of the North Atlantic circulation, which suggests that solar insolation also exerts a profound influence on the East Asian Monsoon (Chu et al., 2014; Liu et al., 2000; Wang et al., 2016).

4.3. Prediction of future climate change on annual to decadal scales

Since the climate of the past 2000 years provides the background for predicting and simulating future climate changes, both PAGES and IPCC have established a global database comprised of records from various geological archives spanning this interval. To predict and simulate future climate change, these archives need to be comparable with, and potentially combined with, instrumental data and historical archives which have a high temporal resolution. Therefore, research on climate change over the past 2000 years focuses on annual-decadal scales. However, few geological archives and proxies of climate change can achieve annual-decadal resolution. In this respect, maar lakes with annually-laminated sediments can provide seasonal-annual-scale records which enable the study of climate change at a high temporal resolution, especially in low-mid latitude continental regions.

The maar lake records from Hoya La Alberca and Hoya Rincón de Parangueo in Mexico, span the interval 1830–1980 CE, and confirm that El Niño was the cause of regional droughts in Central Mexico that were recorded in historical documents (Kienel et al., 2009). El Niño–Southern Oscillation (ENSO) frequencies (Allan et al., 2003) of ~2–8 years are also evident in an annually-laminated sediment sequence spanning the past 530 years from Maar Lake Twintaung in Myanmar, which confirms the close relationship between regional rainfall and ENSO (Sun et al., 2016b). The ENSO frequencies not only occur in modern or Quaternary climatic records, but they are also present in Miocene sedimentary sequences (Lindqvist and Lee, 2009). Periodic variations (~3–7 years) reflected in the couplet thickness of laminated diatomite from Foulden Maar in New Zealand, which is comparable with Quaternary records of ENSO variability and indicates that the climate at low altitudes in New Zealand in the beginning of the Early Miocene was seasonal and strongly ocean-influenced, just like today (Lindqvist and Lee, 2009). In addition to ENSO, Schwabe sunspot cycles are also clearly resolved on an annual-decadal timescale in maar lake sedimentary sequences. The 11-year cycle detected in annually laminated sediments from Holzmaar in Germany and Lake Twintaung in Myanmar provide strong evidence for the presence of sunspot cycles in Quaternary sediments (Vos et al., 1997; Sun et al., 2016b). Furthermore, the 5.5-year cycle found in the laminated Eocene sediments of Eckfeld Maar from the Eifel region in Germany and the 11-year cycle

found in Miocene laminated sediments of Foulden Maar in New Zealand support the conclusion that a sun-climate relationship at the annual-decadal scale existed long before the Quaternary (Mingram, 1998; Lindqvist and Lee, 2009).

In China, periodicities of 2.6, 3.5, 53–55, 87–89 and 105–110 years were detected in a high-resolution minor element dataset covering the past 1300 years from the annually-laminated sediments of Lake Xiaolongwan, in Northeast China (Chu et al., 2013). The occurrence of classical ENSO periodicities and solar oscillations in the record implies that the variability of precipitation in Northeast China was associated with ENSO and solar activity during the past 1300 years (Chu et al., 2013).

5. Quantitative paleotemperature reconstructions

Numerous proxies have been used to produce climatic reconstructions from maar sediments, including analysis of the stable isotopes of oxygen and carbon, magnetic properties, palynology, and aquatic biological indicators such as diatoms, chrysophytes, chironomids and ostracods (Marchetto et al., 2015). Some of these indicators are still in the process of development for application in quantitative palaeoclimatic reconstruction. Traditionally, quantitative reconstructions are based on the modern analogue approach in which paleodata are compared with a modern database. The most successful cases are reconstructions of temperature and precipitation using pollen records (Nakagawa et al., 2002). However, considering the time lag (150–250 years) between the responses of vegetation to climate change (Chapin and Starfield, 1997), new climatically-sensitive proxies that can be measured at high temporal resolution are needed.

The most promising among such new proxies are long-chain alkenones (LCAs) and glycerol dialkyl glycerol tetraethers (GDGTs). They have been widely used for quantitative paleotemperature reconstruction, since they respond rapidly to temperature change during the growing season. LCAs are temperature-dependent lipids produced by several species of haptophyceae that are widely distributed in marine and lacustrine sediments (Sun et al., 2007). Previous studies indicated that the $U^{k_{37}}/U^{k_{37}}$ index, originally developed using di-, tri- and tetra-unsaturated ketones ($C_{37:2}$, $C_{37:3}$, and $C_{37:4}$), was linearly related to sea-surface temperature or culture temperature within the range of roughly 10–22 °C (Brassell et al., 1986; Sun et al., 2007). LCAs are widely applied in sea surface temperature reconstruction, and they are now attracting increasing attention for lake temperature reconstruction, especially in volcanic lakes.

Glycerol dialkyl glycerol tetraethers (GDGTs) are a group of ubiquitous membrane lipids produced by archaea or bacteria, which consist of two types of molecule with different structures and sources, named isoprenoid GDGTs (iGDGTs) and branched GDGTs (brGDGTs). These compounds mainly occur in soils and lake sediments. The development of the Methylation index of Branched Tetraethers (MBT), the Cyclisation ratio of Branched Tetraethers (CBT) (Weijers et al., 2007b) and brGDGTs-based temperature calibration (Foster et al., 2016; Loomis et al., 2012; Pearson et al., 2011; Tierney et al., 2010; Weijers et al., 2007b; Zink et al., 2016), have enabled the reconstruction of mean annual air temperature (MAAT) variations and have generated significant interest within the terrestrial palaeoclimate research community working on lake sediments (Castañeda et al., 2011; Chu et al., 2017; Fawcett et al., 2011; Hu et al., 2015; Zink et al., 2010), peat deposits (Wang et al., 2017; Zheng et al., 2017), loess (Peterse et al., 2014; Peterse et al., 2011) and other sediments (Weijers et al., 2007a). The MBT and CBT proxies are defined and their relationship between the distribution of brGDGTs and mean annual air temperature in numerous studies of soil brGDGTs worldwide are discussed in Weijers et al. (2007b). The GDGTs-temperature calibration was built by analyzing the relationship between GDGTs from many lakes sites worldwide and mean annual air temperature, and temperature reconstructions from the GDGTs-

Table 3

Location, hydrological status and genetic classification of maar lakes in which GDGTs and LCA are detected.

No.	Lake	Latitude	Longitude	Elevation (m)	Water depth (m)	pH	Volcanic area	Country	Sample type	Biomarkers	Reference
1	Grande di Monticchio	40°56'N	15°36'E	656	36	6.7	Monte Vulture	Italy	Surface	GDGTs	Blaga et al., 2010
2	Nyamiteza	0°26'N	30°14'E	1254	34	8.2	Kyatwa	Uganda	Surface	GDGTs	Tierney et al., 2010
3	Kanyamukali	0°24'N	30°14'E	1161	10	8.1					
4	Lugembe	0°27'N	30°17'E	1280	18	7.3					
5	Nyabikere	0°30'N	30°20'E	1463	45	6.8					
6	Ntambi	0°25'N	30°14'E	1158	42	9.3					
7	Donglongwan	42°26'N	126°31'E	620	127	8.4	Longgang	China	Surface	GDGTs	Chu et al., 2005b;
8	Sihailongwan	42°17'N	126°36'E	778	50	7.9			Surface, core	GDGTs, LCA	Sun et al., 2011;
9	Sanjiaolongwan	42°22'N	126°26'E	722	76	7.8			Surface	LCA	Hu et al., 2015, 2016
10	Xiaolongwan	42°18'N	126°21'E	643	15	6.8			Surface	GDGTs	
11	Huguangyan	21°09'N	110°17'E	16	22	7.7	Leiqiong		Surface, trap, core	GDGTs, LCA	
12	Pupuke	36°47'S	174°46'E	12	57	7.7	Auckland	New Zealand	Core	GDGTs	Heyng et al., 2015

temperature calibration of lake sediments has been successfully applied in different lakes.

There is an increasing use of LCAs and GDGTs in lacustrine sediments for terrestrial environmental reconstruction, including the sediments of maar lakes. LCAs have been found in >100 lakes worldwide, including in three maar lakes in China: Sihailongwan and Sanjiaolongwan in Northeast China and Huguangyan in South China (Chu et al., 2005b). GDGTs have been found in over 300 lakes globally, 11 of which are maar lakes, including one in Italy (Grande di Monticchio, Blaga et al., 2010), five in Uganda (Nyamiteza, Kanyamukali, Lugembe, Nyabikere and Ntambi, Tierney et al., 2010), four in China (Donglongwan, Sihailongwan, Xiaolongwan and Huguangyan, Chu et al., 2005b; Hu et al., 2015, 2016; Sun et al., 2011) and one in New Zealand (Pupuke, Heyng et al., 2015). Table 3 lists the pH values of LCAs and GDGTs-containing maar lakes, and the values vary from 6.6–9.3, from neutral to alkaline. This neutral regions in pH is most likely caused by the pH buffers of $\text{HSO}_4^-/\text{SO}_4^{2-}$ and $\text{H}_2\text{CO}_3/\text{HCO}_3^-$ (Marini et al., 2003), and several volcanic lakes with carbonate inputs may have pH values > 8 (Christenson et al., 2015; Pecoraino et al., 2015). This finding may indicate that carbonate-rich maar lakes have higher pH values and tend to contain abundant LCAs and GDGTs, which

accords with the previous conclusion that sulfate-dominated or carbonate-dominated lakes also affect the occurrence and abundance of LCAs (Sun et al., 2004; Toney et al., 2010). The verification of this hypothesis requires further work, especially on unexplored maar lakes.

The high-resolution GDGTs-based temperature records from the sediments of Maar Lake Huguangyan in tropical China reveal that the mean temperature during the Oldest Dryas and Younger Dryas cold events, were 17.8 °C and 19.8 °C, respectively; these values are 5.6 °C and 3.6 °C colder than the respective modern mean annual air temperatures (Chu et al., 2017, Fig. 4). During the Holocene, the mean temperature of the Southern Hemisphere was lower in the early Holocene (15.6 °C), and it continued to rise until 1600 years cal. BP (18.4 °C), which is revealed by a GDGTs-based temperature record from Lake Pupuke, New Zealand (Heyng et al., 2015, Fig. 4). The high-resolution LCAs-based reconstruction of the growing season temperature over the past 1600 years at Lake Sihailongwan, in Northeast China, indicates that the most pronounced cold spells occurred during 480–860 CE, 1260–1300 CE, 1510–1570 CE and 1800–1900 CE, with a temperature decrease of about 1 °C compared to today (Chu et al., 2011, Fig. 4). As more records of LCAs and GDGTs from maar lakes are obtained, high-




Fig. 4. Paleotemperature reconstructions from maar and maar lakes and comparison with the temperature and $\delta^{18}\text{O}$ record of the Greenland ice core. Black curve: winter temperature reconstruction from Huguangyan Maar Lake in China (Chu et al., 2017); green curve: annual mean temperature reconstruction from Pupuke Maar Lake in New Zealand (Heyng et al., 2015); dark blue curve: annual mean air temperature reconstruction from Huguangyan Maar Lake in China (Hu et al., 2015); red curve: growing season temperature reconstruction from Sihailongwan Maar Lake in China (Chu et al., 2011); purple curve: Greenland temperature during the Last Deglaciation (Buizert et al., 2014); light blue curve: $\delta^{18}\text{O}$ record of the Greenland NGRIP ice core (Rasmussen et al., 2006, 2008). The dark shading represents cold events such as the Younger Dryas (YD) and the Oldest Dryas (OD). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

resolution quantitative temperature reconstructions at the regional and continental scales will become possible (Fig. 4).

6. Summary

Palaeoclimatic records of lacustrine sediments from >40 maars have been reported globally, providing continuous records spanning > 400 kyr. In China, numerous palaeoclimatic records from maar sediments have been reported from Leiqiong Volcanic Field and Longgang Volcanic Field.

Absolute varve chronologies spanning over 14,000 years have been established from maar lakes in Longgang Volcanic Field, and tephrochronology studies indicate that the AT tephra and the B-Tm tephra layers can be used as isochrones in East Asia. These results imply that the varve-based ages of maar sediments in Longgang Volcanic Field can be transferred to other sequences where the AT tephra and the B-Tm tephra are present, enabling the direct correlation of the maar sediment sequences in Northeast China and Japan, as well as with ice core sequences in Greenland.

Tianyang Maar from Leiqiong Volcanic Field provides a continuous sedimentary record spanning at least 400 kyr, and the record confirms that vegetation and climatic history of low-latitude tropical China was controlled by the ~10 kyr glacial-interglacial cycle. At a millennial scale, D-O cycles are present in the records of maar lakes from both low and mid latitude China. In particular, typical ENSO periodicities and evidence of solar oscillations can be identified in annual-scale minor element records from maar sediments in Longgang Volcanic Field, which implies that precipitation variability in Northeast China was related to ENSO and solar activity over the past 1300 years.

In addition to conventional climatic proxies, several new quantitative proxies, including LCAs and GDGTs, are being explored. These new proxies have already delivered high-resolution temperature records since the last deglaciation in China.

Maar sediments from Leiqiong and Longgang Volcanic Fields in China, whose precise chronologies are based on varves, isotope dating and tephrochronology, provide valuable palaeoclimatic records for this key region of East Asian Monsoon influence, and they contribute significantly to the global paleoenvironmental and palaeoclimatic database for maars and maar lakes.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (grants No. 41320104006, 41561144010 and 41572353), the Key Research Project of Frontier Sciences of Chinese Academy of Sciences (QYDZ-SSW-DQC030), and the Chinese Postdoctoral Foundation (No. BX201600160, 2015LH001, and 2017M620897). We thank Dr. Karoly Németh from Massey University, New Zealand; Dr. Dmitri Rouwet from Istituto Nazionale di Geofisica e Vulcanologia, Italy; Dr. Xavier Bolós from the Institute of Geophysics, Universidad Nacional Autónoma De México; and Dr. Martin Koziol from Maarmuseum Manderscheid, Germany, for their help in collecting information on maar volcanoes and for improving the quality of the manuscript.

References

- Allan, R.J., Reason, C.J.C., Lindesay, J.A., Ansell, T.J., 2003. 'Protracted' ENSO episodes and their impacts in the Indian Ocean region. *Deep-Sea Res. II Top. Stud. Oceanogr.* 50 (12), 2331–2347.
- Ariztegui, D., Chondrogianni, C., Lami, A., Guilizzoni, P., Lafargue, E., 2001. Lacustrine organic matter and the Holocene paleoenvironmental record of Lake Albano central Italy. *J. Paleolimnol.* 26 (3), 283–292.
- Augustinus, P., 2007. NZ-Maars; extracting high resolution paleoclimate records from maar crater lakes, Auckland, New Zealand. *Past Global Changes*. 15, pp. 18–20.
- Augustinus, P., 2012. The Last Glacial cycle viewed from Onepoto and Pukaki Maar Lakes, Auckland, New Zealand. *Quat. Int.* 279, 28.
- Augustinus, P., Cochran, U., Kattel, G., D'Costa, D., Shane, P., 2012. Late Quaternary paleolimnology of Onepoto maar, Auckland, New Zealand: Implications for the drivers of regional paleoclimate. *Quat. Int.* 253, 18–31.
- Bannister, J.M., Conran, J.G., Lee, D.E., 2012. Lauraceae from rainforest surrounding an early Miocene maar lake, Otago, southern New Zealand. *Rev. Palaeobot. Palynol.* 178, 13–34.
- Barker, P., Williamson, D., Gasse, F., Gibert, E., 2003. Climatic and volcanic forcing revealed in a 50,000-year diatom record from Lake Massoko, Tanzania. *Quat. Res.* 60 (3), 368–376.
- Baumann, K.H., Lackschewitz, K.S., Mangerud, J., Spielhagen, R.F., Wolf-Welling, T.C.W., Henrich, R., Kassens, H., 1995. Reflection of Scandinavian ice sheet fluctuations in Norwegian Sea sediments during the past 150,000 years. *Quat. Res.* 43 (2), 185–197.
- Berger, A., 1988. Milankovitch theory and climate. *Rev. Geophys.* 26 (4), 624–657.
- Berger, A., Loutre, M.F., 2002. An exceptionally long interglacial ahead? *Science* 297 (5585), 1287–1288.
- Blaga, C.I., Reichert, G.J., Schouten, S., Lotter, A.F., Werne, J.P., Kosten, S., Mazzeo, N., Lacerot, G., Damsté, J.S.S., 2010. Branched glycerol dialkyl glycerol tetraethers in lake sediments: can they be used as temperature and pH proxies? *Org. Geochem.* 41 (11), 1225–1234.
- Bond, G.C., Lotti, R., 1995. Iceberg discharges into the North Atlantic on millennial time scales during the last glaciation. *Science* 267 (5200), 1005.
- Bond, G., Heinrich, H., Broecker, W., Labeyrie, L., McManus, J., Andrews, J., Huon, S., Jantschik, R., Clasen, S., Simet, C., 1992. Evidence for massive discharges of icebergs into the North Atlantic ocean during the last glacial period. *Nature* 360 (6401), 245–249.
- Bond, G., Showers, W., Cheseby, M., Lotti, R., Almasi, P., de Menocal, P., Priore, P., Cullen, H., Hajdas, I., Bonani, G., 1997. A pervasive millennial-scale cycle in North Atlantic Holocene and Glacial Climates. *Science* 278 (5341), 1257–1266.
- Bond, G., Kromer, B., Beer, J., Muscheler, R., Evans, M.N., Showers, W., Hoffmann, S., Lotti-Bond, R., Hajdas, I., Bonani, G., 2001. Persistent solar influence on North Atlantic climate during the Holocene. *Science* 294 (5549), 2130–2136.
- Brassell, S., Eglinton, G., Marlowe, I.T., Pflaumann, U., Sarinthein, M., 1986. Molecular stratigraphy: a new tool for climatic assessment. *Nature* 320 (6058), 129–133.
- Brauer, A., Endres, C., Negendank, J.F.W., 1999. Lateglacial calendar year chronology based on annually laminated sediments from Lake Meerfelder Maar, Germany. *Quat. Int.* 61 (1), 17–25.
- Broecker, W.S., 1966. Absolute dating and the astronomical theory of glaciation. *Science* 151 (3708), 299–304.
- Brohan, P., Kennedy, J.J., Harris, I., Tett, S.F.B., Jones, P.D., 2006. Uncertainty estimates in regional and global observed temperature changes: a new data set from 1850. *J. Geophys. Res.* 111, D12106.
- Büchel, G., 1993. Maars of the Westeifel, Germany. *Paleolimnology of European Maar Lakes*. pp. 1–13.
- Büchel, G., Lorenz, V., 1982. Zum Alter des Maarvulkanismus der Westeifel. *Neues Jahrbuch fuer Geologie, Paleontologie. Fortschr. Mineral.* 163, 1–22.
- Buizert, C., Gkinis, V., Severinghaus, J.P., He, F., Lecavalier, B.S., Kindler, P., Leuenberger, M., Carlson, A.E., Vinther, B., Masson-Delmotte, V., White, J.W.C., Liu, Z., Otto-Bliesner, B., Brook, E.J., 2014. Greenland temperature response to climate forcing during the last deglaciation. *Science* 345 (6201), 1177.
- Castañeda, I.S., Werne, J.P., Johnson, T.C., Powers, L.A., 2011. Organic geochemical records from Lake Malawi (East Africa) of the last 700 years, part II: biomarker evidence for recent changes in primary productivity. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 303 (1–4), 140–154.
- Chapin, F.S., Starfield, A.M., 1997. Time lags and novel ecosystems in response to transient climatic change in Arctic Alaska. *Clim. Chang.* 35 (4), 449–461.
- Chaparro, M.A.E., Böhnelt, H.N., Byrne, R., Nowaczyk, N.R., Molina-Garza, R.S., Park, J., Negendank, J.F.W., 2008. Palaeomagnetic secular variation and rock-magnetic studies of Holocene sediments from a maar lake (Hoya de San Nicolas) in Central Mexico. *Geophys. J. Int.* 175 (2), 462–476.
- Christenson, B., Németh, K., Rouwet, D., Tassi, F., Vandemeulebrouck, J., Varekamp, J.C., 2015. Volcanic lakes. In: Rouwet, D., Christenson, B., Tassi, F., Vandemeulebrouck, J. (Eds.), *Volcanic Lakes*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 1–20.
- Chu, G., Liu, J., Liu, D., 2000. Discrimination of two kinds of sedimentary laminae in maar lakes of China. *Chin. Sci. Bull.* 45 (24), 2292–2295.
- Chu, G., Liu, J., Sun, Q., Lu, H., Gu, Z., Wang, W., Liu, T., 2002. The "Mediaeval Warm Period" drought recorded in Lake Huguangyan, tropical South China. *The Holocene* 12 (5), 511–516.
- Chu, G., Liu, J., Schettler, G., Li, J., Sun, Q., Gu, Z., Lu, H., Liu, Q., Liu, T., 2005a. Sediment fluxes and varve formation in Sihailongwan, a maar lake from northeastern China. *J. Paleolimnol.* 34 (3), 311–324.
- Chu, G., Sun, Q., Li, S., Zheng, M., Jia, X., Lu, C., Liu, J., Liu, T., 2005b. Long-chain alkenone distributions and temperature dependence in lacustrine surface sediments from China. *Geochim. Cosmochim. Acta* 69 (21), 4985–5003.
- Chu, G., Sun, Q., Rioual, P., Boltovskoy, A., Liu, Q., Sun, P., Han, J., Liu, J., 2008. Dinocyst microlaminations and freshwater "red tides" recorded in Lake Xiaolongwan, northeastern China. *J. Paleolimnol.* 39 (3), 319–333.
- Chu, G., Sun, Q., Wang, X., Li, D., Rioual, P., Qiang, L., Han, J., Liu, J., 2009. A 1600 year multiproxy record of paleoclimatic change from varved sediments in Lake Xiaolongwan, northeastern China. *J. Geophys. Res.* 114 (D22), D22108.
- Chu, G., Sun, Q., Wang, X., Liu, M., Lin, Y., Xie, M., Shang, W., Liu, J., 2011. Seasonal temperature variability during the past 1600 years recorded in historical documents and varved lake sediment profiles from northeastern China. *The Holocene* 22 (7), 785–792.
- Chu, G., Sun, Q., Li, S., Lin, Y., Wang, X., Xie, M., Shang, W., Li, A., Yang, K., 2013. Minor element variations during the past 1300 years in the varved sediments of Lake Xiaolongwan, north-eastern China. *GFF* (1–8).

- Chu, G., Sun, Q., Xie, M., Lin, Y., Shang, W., Zhu, Q., Shan, Y., Xu, D., Rioual, P., Wang, L., 2014. Holocene cyclic climatic variations and the role of the Pacific Ocean as recorded in varved sediments from northeastern China. *Quat. Sci. Rev.* 102, 85–95.
- Chu, G., Sun, Q., Zhu, Q., Shan, Y., Shang, W., Ling, Y., Su, Y., Xie, M., Wang, X., Liu, J., 2017. The role of the Asian winter monsoon in the rapid propagation of abrupt climate changes during the last deglaciation. *Quat. Sci. Rev.* 177, 120–129.
- Conran, J.G., Bannister, J.M., Mildenhall, D.C., Lee, D.E., Chacon, J., Renner, S.S., 2014. Leaf fossils of *Luzuriaga* and a monocot flower with in situ pollen of *Liliacites contortus* Mildenh. & Bannister sp. nov. (*Alstroemeriaceae*) from the Early Miocene. *Am. J. Bot.* 101 (1), 141–155.
- Creer, K.M., 1996. The EUROMAARS Project: A Multidisciplinary Approach Directed to the Reconstruction of Palaeoenvironmental, Palaeoclimatic and Palaeogeomagnetic Changes Recorded by Sediments Deposited in Maar Lakes Located in Western Europe. Pergamon.
- Creer, K., Morris, A., 1996. Proxy-climate and geomagnetic palaeointensity records extending back to ca. 75,000 BP derived from sediments core from Lago Grande Di Monticchio, southern Italy. *Quat. Sci. Rev.* 15 (2–3), 167–188.
- Damon, P.E., Jirakovic, J.L., 1992. The sun as a low-frequency harmonic oscillator. *Radio-carbon* 34 (2), 199–205.
- Dansgaard, W., Johnsen, S.J., Clausen, H.B., Dahl-Jensen, D., Gundestrup, N.S., Hammer, C.U., Hvidberg, C.S., Steffensen, J.P., Sveinbjörnsdóttir, A.E., Jouzel, J., Bond, G., 1993. Evidence for general instability of past climate from a 250-kyr ice-core record. *Nature* 364, 218.
- D'Costa, D.M., Kershaw, A.P., 1995. A Late Pleistocene and Holocene pollen record from Lake Terang, Western Plains of Victoria, Australia. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 113 (1), 57–67.
- De Geer, G., 1912. A geochronology of the last 12000 years. *Compte Rendu 11 Congress Geologic International, 11th International Geological Congress (1910)*, Stockholm, Sweden, pp. 241–253.
- De Soler, B.G., Vall-Lloera, G.C., Van der Made, J., Oms, O., Agustí, J., Sala, R., Blain, H.A., Burjachs, F., Claude, J., Catalan, S.G., Riba, D., Rosillo, R., 2012. A new key locality for the Pliocene vertebrate record of Europe: the Camp dels Ninots maar (NE Spain). *Geol. Acta* 10 (1), 1–17.
- Dimitriadis, S., Cranston, P.S., 2001. An Australian Holocene climate reconstruction using *Chironomidae* from a tropical volcanic maar lake. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 176 (1–4), 109–131.
- Disnar, J.-R., Stefanova, M., Bourdon, S., Laggoun-Défarge, F., 2005. Sequential fatty acid analysis of a peat core covering the last two millennia (Tritivakely lake, Madagascar): diagenesis appraisal and consequences for palaeoenvironmental reconstruction. *Org. Geochem.* 36 (10), 1391–1404.
- Fan, Q., Liu, R., Wei, H., Sui, J., Li, N., 1999. The petrology and geochemistry of Jinlongdingzi modern active volcano in Longgang area. *Acta Petrol. Sin.* 15 (4), 584–589 (in Chinese with English abstract).
- Fan, Q., Sui, J., Liu, R., Wei, H., Li, D., Sun, Q., Li, N., 2002. Periods of Quaternary volcanic activity in Longgang area, Jilin province. *Acta Petrol. Sin.* 18 (4), 495–500 (in Chinese with English abstract).
- Fawcett, P.J., Werne, J.P., Anderson, R.S., Heikoop, J.M., Brown, E.T., Berke, M.A., Smith, S.J., Goff, F., Donohoo-Hurley, L., Cisneros-Dozal, L.M., Schouten, S., Sinninghe Damste, J.S., Huang, Y., Toney, J., Fessenden, J., Woldegabriel, G., Atudorei, V., Geissman, J.W., Allen, C.D., 2011. Extended megadroughts in the southwestern United States during Pleistocene interglacials. *Nature* 470 (7335), 518–521.
- Follieri, M., Giardini, M., Magri, D., Sadori, L., 1998. Palynostratigraphy of the Last Glacial Period in the Volcanic Region of Central Italy. *Quat. Int.* 47–48, 3–20.
- Foster, L.C., Pearson, E.J., Juggins, S., Hodgson, D.A., Saunders, K.M., Verleyen, E., Roberts, S.J., 2016. Development of a regional glycerol dialkyl glycerol tetraether (GDGT)-temperature calibration for Antarctic and sub-Antarctic lakes. *Earth Planet. Sci. Lett.* 433, 370–379.
- Fox, B., Wilson, G.S., Lee, D.E., Gorman, A.R., Kaulfuss, U., Jones, D., 2010. Climate and environment in early Miocene New Zealand: new evidence from an Otago maar lake. *Geoscience Society of New Zealand Miscellaneous Publication*. 129A, p. 98.
- Fox, B., Wilson, G.S., Lee, D.E., Haworth, M., Wartho, J., Bannister, J., Kaulfuss, U., Clark, K.J., Litchfield, N.J., 2011. Causes and consequences of climate change during the Mi-1 event: paleoclimate data from the Foulden Maar core. *Geoscience Society of New Zealand Miscellaneous Publication*. 130A, p. 39.
- Frank, A.H.E., 1969. Pollen stratigraphy of the lake of vico (Central Italy). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 6, 67–85.
- Fuhrmann, A., Mingram, J., Lücke, A., Lu, H., Horsfield, B., Liu, J., Negendank, J.F.W., Schleser, G.H., Wilkes, H., 2003. Variations in organic matter composition in sediments from Lake Huguang Maar (Huguangyan), south China during the last 68 ka: implications for environmental and climatic change. *Org. Geochem.* 34 (11), 1497–1515.
- Ganopolski, A., Winkelmann, R., Schellnhuber, H.J., 2016. Critical insulation- CO_2 relation for diagnosing past and future glacial inception. *Nature* 529 (7585), 200–203.
- Garcin, Y., Williamson, D., Taieb, M., Vincens, A., Mathé, P.-E., Majule, A., 2006. Centennial to millennial changes in maar-lake deposition during the last 45,000 years in tropical Southern Africa (Lake Masoko, Tanzania). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 239 (3–4), 334–354.
- Gibert, E., Bergonzini, L., Massault, M., Williamson, D., 2002. AMS- ^{14}C chronology of 40.0 cal ka BP continuous deposits from a crater lake (Lake Masoko, Tanzania): modern water balance and environmental implications. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 187 (3–4), 307–322.
- Giguet-Covex, C., Arnaud, F., Poulenard, J., Enters, D., Reyss, J.-L., Millet, L., Lazzarotto, J., Vidal, O., 2010. Sedimentological and geochemical records of past trophic state and hypolimnetic anoxia in large, hard-water Lake Bourget, French Alps. *J. Paleolimnol.* 43 (1), 171–190.
- Giresse, P., Maley, J., Brenac, P., 1994. Late Quaternary palaeoenvironments in the Lake Barombi Mbo (West Cameroon) deduced from pollen and carbon isotopes of organic matter. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 107 (1–2), 65–78.
- Goth, K., Suhr, P., 2003. Algal dominated sediments in maar lakes. *Foldtani Kutatas* 40 (1–2), 22–24.
- Graettinger, A.H., 2018. Trends in maar crater size and shape using the global Maar Volcano Location and Shape (MaarVLS) database. *J. Volcanol. Geotherm. Res.* 357, 1–13.
- Grein, M., Utescher, T., Wilde, V., Roth-Nebelsick, A., 2011. Reconstruction of the middle Eocene climate of Messel using palaeobotanical data. *Neues Jahrbuch Fur Geologie Und Paläontologie-Abhandlungen* 260 (3), 305–318.
- Grootes, P.M., Stuiver, M., White, J.W.C., Johnsen, S., Jouzel, J., 1993. Comparison of oxygen isotope records from the GISP2 and GRIP Greenland ice cores. *Nature* 366 (6455), 552–554.
- Hajdas, I., Zolitschka, B., Ivy-Ochs, S.D., Beer, J., Bonani, G., Leroy, S.A., Negendank, J.W., Ramrath, M., Suter, M., 1995. AMS radiocarbon dating of annually laminated sediments from Lake Holzmaar, Germany. *Quat. Sci. Rev.* 14 (2), 137–143.
- Hao, Q., Wang, L., Oldfield, F., Peng, S., Qin, L., Song, Y., Xu, B., Qiao, Y., Bloemendal, J., Guo, Z., 2012. Delayed build-up of Arctic ice sheets during 400,000-year minima in insolation variability. *Nature* 490 (7420), 393–396.
- Heyng, A.M., Mayr, C., Lücke, A., Moschen, R., Wissel, H., Striewski, B., Bauersachs, T., 2015. Middle and Late Holocene paleotemperatures reconstructed from oxygen isotopes and GDGTs of sediments from Lake Pupuke, New Zealand. *Quat. Int.* 374, 3–14.
- Hunag, Z., Cai, F., Han, Z., 1993. Quaternary Volcanoes of Leizhou Peninsula and Northern Hainan Island. Science Press, Beijing (in Chinese).
- Hu, J., Zhou, H., Peng, P.A., Yang, X., Spiro, B., Jia, G., Wei, G., Ouyang, T., 2015. Reconstruction of a paleotemperature record from 0.3–3.7 ka for subtropical South China using lacustrine branched GDGTs from Huguangyan Maar. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 435 (Supplement C), 167–176.
- Hu, J., Zhou, H., Peng, P.A., Spiro, B., 2016. Seasonal variability in concentrations and fluxes of glycerol dialkyl glycerol tetraethers in Huguangyan Maar Lake, SE China: implications for the applicability of the MBT-CBT paleotemperature proxy in lacustrine settings. *Chem. Geol.* 420 (Supplement C), 200–212.
- Huang, X.L., Xu, Y.G., Lo, C.H., Wang, R.C., Lin, C.Y., 2007. Exsolution lamellae in a clinopyroxene megacryst aggregate from Cenozoic basalt, Leizhou Peninsula, South China: petrography and chemical evolution. *Contrib. Mineral. Petrol.* 154 (6), 691–705.
- Jones, R.N., McMahon, T.A., Bowler, J.M., 2001. Modelling historical lake levels and recent climate change at three closed lakes, Western Victoria, Australia (c.1840–1990). *J. Hydrol.* 246 (1), 159–180.
- Juvigné, E., Camus, G., de Herve, A.D.G., 1993. Maars of northern Auvergne (Massif Central, France): state of knowledge. In: Negendank, J.F.W., Zolitschka, B. (Eds.), *Paleolimnology of European Maar Lakes*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 81–94.
- Kattel, G., Sirocko, F., 2011. Palaeocladocera as indicators of environmental, cultural and archaeological developments in Eifel maar lakes region (West Germany) during the Lateglacial and Holocene periods. *Hydrobiologia* 676 (1), 203–221.
- Kaulfuss, U., Wappler, T., Heiss, E., Larivière, M.C., 2011. Aneurys sp from the early Miocene Foulden Maar, New Zealand: the first Southern Hemisphere record of fossil Aradidae (Insecta: Hemiptera: Heteroptera). *J. R. Soc. N. Z.* 41 (4), 279–285.
- Kienel, N., Bowen, S.W., Byrne, R., Park, J., Böhnelt, H., Dulski, P., Luhr, J.F., Siebert, L., Haug, G.H., Negendank, J.F., 2009. First lacustrine varve chronologies from Mexico: impact of droughts, ENSO and human activity since AD 1840 as recorded in maar sediments from Valle de Santiago. *J. Paleolimnol.* 42 (4), 587–609.
- Kliem, P., Enters, D., Hahn, A., Ohlendorf, C., Lisé-Pronovost, A., St-Onge, G., Wastegård, S., Zolitschka, B., 2013. Lithology, radiocarbon chronology and sedimentological interpretation of the lacustrine record from Laguna Potrok Aike, southern Patagonia. *Quat. Sci. Rev.* 71 (71), 54–69.
- Lamb, A.L., Leng, M.J., Lamb, H.F., Mohammed, M.U., 2000. A 9000-year oxygen and carbon isotope record of hydrological change in a small Ethiopian crater lake. *The Holocene* 10 (2), 167–177.
- Lami, A., Niessen, F., Guilizzoni, P., Masferro, J., Belis, C.A., 1994. Palaeolimnological studies of the eutrophication of volcanic Lake Albano (Central Italy). *J. Paleolimnol.* 10 (3), 181–197.
- Lane, C., Blockley, S., Lotter, A., Finsinger, W., Filippi, M., Matthews, I., 2012. A regional tephrostratigraphic framework for central and southern European climate archives during the Last Glacial to Interglacial transition: comparisons north and south of the Alps. *Quat. Sci. Rev.* 36, 50–58.
- Lane, C.S., Brauer, A., Blockley, S.P., Dulski, P., 2013. Volcanic ash reveals time-transgressive abrupt climate change during the Younger Dryas. *Geology* 41 (12), 1251–1254.
- Lane, C.S., Brauer, A., Martín-Puertas, C., Blockley, S.P., Smith, V.C., Tomlinson, E.L., 2015. The Late Quaternary tephrostratigraphy of annually laminated sediments from Meerfelder Maar, Germany. *Quat. Sci. Rev.* 122, 192–206.
- Lea, D.W., Pak, D.K., Spero, H.J., 2000. Climate impact of Late Quaternary equatorial Pacific sea surface temperature variations. *Science* 289 (5485), 1719–1724.
- Lenz, O.K., Wilde, V., Riegel, W., Harns, F.J., 2010. A 600 k.y. record of El Niño-Southern Oscillation (ENSO): evidence for persisting teleconnections during the Middle Eocene greenhouse climate of Central Europe. *Geology* 38 (7), 627–630.
- Lenz, O.K., Wilde, V., Riegel, W., 2011. Short-term fluctuations in vegetation and phytoplankton during the Middle Eocene greenhouse climate: a 640-kyr record from the Messel oil shale (Germany). *Int. J. Earth Sci.* 100 (8), 1851–1874.
- Lenz, O.K., Wilde, V., Riegel, W., 2014. Palynology as a high-resolution tool for cyclostratigraphy in Middle Eocene lacustrine sediments: the outstanding record of Messel (Germany). In: Rocha, R., Pais, J., Kullberg, J.C., Finney, S. (Eds.), *Strati 2013*. Springer Geology, pp. 113–117.
- Lenz, O.K., Wilde, V., Riegel, W., 2017. ENSO- and solar-driven sub-Milankovitch cyclicity in the Palaeocene greenhouse world: high-resolution pollen records from Eocene Lake Messel, Germany. *J. Geol. Soc.* 174 (1), 110–128.

- Lindqvist, J.K., Lee, D.E., 2009. High-frequency paleoclimate signals from Foulden Maar, Waipiaata Volcanic Field, southern New Zealand: an Early Miocene varved lacustrine diatomite deposit. *Sediment. Geol.* 222 (1), 98–110.
- Lisiecki, L.E., Raymo, M.E., 2009. Diachronous benthic $\delta^{18}\text{O}$ responses during late Pleistocene terminations. *Paleoceanography* 24 (3), PA3210.
- Litt, T., Stebich, M., 1999. Bio- and chronostratigraphy of the lateglacial in the Eifel region, Germany. *Quat. Int.* 61 (1), 5–16.
- Litt, T., Brauer, A., Goslar, T., Merkt, J., Bałaga, K., Müller, H., Ralska-Jasiewiczowa, M., Stebich, M., Negendank, J.F., 2001. Correlation and synchronisation of Lateglacial continental sequences in northern central Europe based on annually laminated lacustrine sediments. *Quat. Sci. Rev.* 20 (11), 1233–1249.
- Liu, J., Liu, T., Chu, G., Xiao, J., Luo, X., 1996. Maar lake and varve chronology. *Quat. Sci.* 16 (4), 353–358 (in Chinese with English abstract).
- Liu, J., Houyuan, L., Negendank, J., Mingram, J., Luo, X., Wang, W., Chu, G., 2000. Periodicity of Holocene climatic variations in the Huguangyan Maar Lake. *Chin. Sci. Bull.* 45 (18), 1712–1717.
- Liu, Q., Zhao Yan, G.U., Liu, J.Q., You, H.T., Yuan, L.H., Chu, G.Q., Xiang Lei, Q.I., Negendank, J.F.W., Jens, M., Georg, S., 2005a. Bulk organic carbon isotopic record of Huguangyan Maar Lake, Southeastern China and its paleoclimatic and paleoenvironmental significance since 62 kaBP. *Mar. Geol. Quat. Geol.* 25 (2), 115–126 (in Chinese with English abstract).
- Liu, Q., Liu, J., Chen, X., You, H., Chu, G., Jingtai, H., Mingram, J., Schettler, G., Negendank, J.W., 2005b. Stable carbon isotope record of bulk organic matter from the Sihailongwan Maar Lake, Northeast China during the past 18.5 ka. *Quat. Sci.* 6, 711–721 (in Chinese with English abstract).
- Liu, J., Chu, G., Han, J., Rioual, P., Jiao, W., Wang, K., 2009. Volcanic eruptions in the Longgang volcanic field, northeastern China, during the past 15,000 years. *J. Asian Earth Sci.* 34 (5), 645–654.
- Liu, J., Chu, G., Liu, Q., Wang, L., Rioual, P., Wu, J., Liu, J., You, H., 2013. Progress of palaeoclimatic and palaeoenvironmental studies on maar lakes. *Bull. Mineral. Petrol. Geochim.* 32 (6), 639–652 (in Chinese with English abstract).
- Loomis, S.E., Russell, J.M., Ladd, B., Street-Perrott, F.A., Sinninghe Damsté, J.S., 2012. Calibration and application of the branched GDGT temperature proxy on East African lake sediments. *Earth Planet. Sci. Lett.* 357–358, 277–288.
- Liu, J.Q., Sun, C.Q., You, H.T., 2018. An overview of global tephrochronology. *Sci. Sin. Terrae* 48, 1–29 (in Chinese with English abstract).
- Lorenz, V., 1973. On the formation of maars. *Bull. Volcanol.* 37 (2), 183–204.
- Lorenz, V., 1975. Formation of phreatomagmatic maar-diatreme volcanoes and its relevance to kimberlite diatremes. *Phys. Chem. Earth* (9), 17–27.
- Lorenz, V., 1986. On the growth of maars and diatremes and its relevance to the formation of tuff rings. *Bull. Volcanol.* 48 (5), 265–274.
- Lorenz, V., 2003. Maar-diatreme volcanoes, their formation, and their setting in hard-rock or soft-rock environments. *Geolines* 15, 72–83.
- Lorenz, V., 2007. Syn- and post-eruptive hazards of maar-diatreme volcanoes. *J. Volcanol. Geotherm. Res.* 159 (1–3), 285–312.
- Lowe, D.J., 2011. Tephrochronology and its application: a review. *Quat. Geochronol.* 6 (2), 107–153.
- Machida, H., Arai, F., 1983. Extensive ash falls in and around the sea of Japan from large late quaternary eruptions. *J. Volcanol. Geotherm. Res.* 18 (1), 151–164.
- Magri, D., 1999. Late Quaternary vegetation history at Lagaccione near Lago di Bolsena (central Italy). *Rev. Palaeobot. Palynol.* 106 (3), 171–208.
- Marchetto, A., Ariztegui, D., Brauer, A., Lami, A., Mercuri, A.M., Sadori, L., Vigliotti, L., Wulf, S., Guilizzoni, P., 2015. Volcanic lake sediments as sensitive archives of climate and environmental change. In: Rouwet, D., Christenson, B., Tassi, F., Vandemeulebrouck, J. (Eds.), *Volcanic Lakes*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 379–399.
- Marini, L., Vetuschci Zuccolini, M., Saldi, G., 2003. The bimodal pH distribution of volcanic lake waters. *J. Volcanol. Geotherm. Res.* 121 (1), 83–98.
- Martin-Puertas, C., Tjallingii, R., Bloemsmma, M., Brauer, A., 2017. Varved sediment responses to early Holocene climate and environmental changes in Lake Meerfelder Maar (Germany) obtained from multivariate analyses of micro X-ray fluorescence core scanning data. *J. Quat. Sci.* 32 (3), 427–436.
- Mezger, J.E., Felder, M., Harms, F.-J., 2013. Crystalline rocks in the maar deposits of Messel: key to understand the geometries of the Messel Fault Zone and diatreme and the post-eruptional development of the basin fill. *Zeitschrift Der Deutschen Gesellschaft Fur Geowissenschaften* 164 (4), 639–662.
- Milankovitch, M., 1941. Canon of Insolation and the Ice-age Problem. *Koniglich Serbische Akademie, Beograd*.
- Mingram, J., 1998. Laminated Eocene maar-lake sediments from Eckfeld (Eifel region, Germany) and their short-term periodicities. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 140 (s1–4), 289–305.
- Mingram, J., Nowaczyk, N., Schettler, G., Luo, X., Jiaqi, L., Negendank, J.F.W., 2000. A 78,000 year record of climatic changes from the South China coast-the Huguang maar lake (Huguangyan). *Terra Nostra* 2000 (6), 342–352.
- Mingram, J., Allen, J.R.M., Bruchmann, C., Liu, J., Luo, X., Negendank, J.F.W., Nowaczyk, N., Schettler, G., 2004a. Maar- and crater lakes of the Long Gang Volcanic Field (N.E. China)—overview, laminated sediments, and vegetation history of the last 900 years. *Quat. Int.* 123–125, 135–147.
- Mingram, J., Schettler, G., Nowaczyk, N., Luo, X., Lu, H., Liu, J., Negendank, J.F.W., 2004b. The Huguang maar lake—a high-resolution record of palaeoenvironmental and palaeoclimatic changes over the last 78,000 years from South China. *Quat. Int.* 122 (1), 85–107.
- Mingram, J., Frank, U., You, H., Stebich, M., Liu, J., 2008. A widespread East Asian chronomarker (Aira-TN tephra) found in varved maar lake sediments (Sihailongwan and Erlongwan) of the Longgang Volcanic Field (NE China). 3th International Maar Conference, Malargüe, Argentina.
- Muhs, D.R., Ager, T.A., Josh, B., Platt, B.J., Dean, W.E., 2003. A late Quaternary record of eolian silt deposition in a maar lake, St. Michael Island, western Alaska. *Quat. Res.* 60 (1), 110–122.
- Nakagawa, T., Tarasov, P.E., Nishida, K., Gotanda, K., Yasuda, Y., 2002. Quantitative pollen-based climate reconstruction in central Japan: application to surface and Late Quaternary spectra. *Quat. Sci. Rev.* 21 (18), 2099–2113.
- Nakagawa, T., Gotanda, K., Haraguchi, T., Danhara, T., Yonenobu, H., Brauer, A., Yokoyama, Y., Tada, R., Takemura, K., Staff, R.A., 2012. SG06, a fully continuous and varved sediment core from Lake Suigetsu, Japan: stratigraphy and potential for improving the radiocarbon calibration model and understanding of late Quaternary climate changes. *Quat. Sci. Rev.* 36 (6), 164–176.
- Narcisi, B., 2001. Palaeoenvironmental and palaeoclimatic implications of the Late-Quaternary sediment record of Vico volcanic lake (central Italy). *J. Quat. Sci.* 16 (3), 245–255.
- Negendank, J.F.W., 1984. Die untersuchung der schwerminerale der seesedimente des Meerfelder Maeres und des “Laacher Bimstufes” in den sedimenten des Meerfelder Maeres, des Hinkelsmaeres und der Hitsche. *CFS. Cour. Forschungsinst. Senck.* 65, 41–47.
- Negendank, J.F.W., Zolitschka, B., 1993a. Paleolimnology of European Maar Lakes. Springer, Berlin Heidelberg.
- Negendank, J.F.W., Zolitschka, B., 1993b. Maars and Maar Lakes of the Westeifel Volcanic Field. Springer, Berlin Heidelberg, pp. 61–80.
- Németh, K., Kereszturi, G., 2015. Monogenetic volcanism: personal views and discussion. *Int. J. Earth Sci.* 104, 2131–2146.
- Németh, K., Goth, K., Martin, U., Csillag, G., Suhr, P., 2008. Reconstructing paleoenvironment, eruption mechanism and paleomorphology of the Pliocene Pula maar, (Hungary). *J. Volcanol. Geotherm. Res.* 177 (2), 441–456.
- Nilsson, A., Muscheler, R., Snowball, I., 2011. Multi-proxy identification of the Laschamp geomagnetic field excursion in Lake Pupuke, New Zealand. *Earth Planet. Sci. Lett.* 311 (1), 155–164.
- Nomade, S., Gauthier, A., Guillou, H., Pastre, J.-F., 2010. $^{40}\text{Ar}/^{39}\text{Ar}$ temporal framework for the Alleret maar lacustrine sequence (French Massif-Central): volcanological and paleoclimatic implications. *Quat. Geochronol.* 5 (1), 20–27.
- Okuno, M., Torii, M., Yamada, K., Shinozuka, Y., Danhara, T., Gotanda, K., Yonenobu, H., Yasuda, Y., 2011. Widespread tephra in sediments from lake Ichi-no-Megata in northern Japan: their description, correlation and significance. *Quat. Int.* 246 (1–2), 270–277.
- Ollier, C.D., 1967. Maars their characteristics, varieties and definition. *Bull. Volcanol.* 31 (1), 45–73.
- Ortiz, J.E., Moreno, L., Torres, T., Vegas, J., Ruiz-Zapata, B., García-Cortés, Á., Galán, L., Pérez-González, A., 2013. A 220 ka palaeoenvironmental reconstruction of the Fuentillejo maar lake record (Central Spain) using biomarker analysis. *Org. Geochem.* 55, 85–97.
- Park, J., Byrne, R., Böhnell, H., Garza, R.M., Conserva, M., 2010. Holocene climate change and human impact, central Mexico: a record based on maar lake pollen and sediment chemistry. *Quat. Sci. Rev.* 29 (5–6), 618–632.
- Parplies, J., Lücke, A., Vos, H., Mingram, J., Stebich, M., Radtke, U., Han, J., Schleser, G.H., 2008. Late glacial environment and climate development in northeastern China derived from geochemical and isotopic investigations of the varved sediment record from Lake Sihailongwan (Jilin Province). *J. Paleolimnol.* 40 (1), 471–487.
- Pastre, J.F., Gauthier, A., Nomade, S., Orth, P., Andrieu, A., Goupille, F., Guillou, H., Kunesch, S., Scaillet, S., Renne, P.R., 2007. The Alleret maar (Massif Central, France): a new lacustrine sequence of the early Middle Pleistocene in western Europe. *Compt. Rendus Géosci.* 339 (16), 987–997.
- Pearson, S.J., Juggins, S., Talbot, H.M., Weckström, J., Rosén, P., Ryves, D.B., Roberts, S.J., Schmidt, R., 2011. A lacustrine GDGT-temperature calibration from the Scandinavian Arctic to Antarctica: renewed potential for the application of GDGT-paleothermometry in lakes. *Geochim. Cosmochim. Acta* 75 (20), 6225–6238.
- Pecoraino, G., D'Alessandro, W., Inguaggiato, S., 2015. The Other Side of the Coin: Geochemistry of Alkaline Lakes in Volcanic Areas. Springer, Berlin Heidelberg, pp. 219–237.
- Peterse, F., Prins, M.A., Beets, C.J., Troelstra, S.R., Zheng, H., Gu, Z., Schouten, S., Damsté, J.S.S., 2011. Decoupled warming and monsoon precipitation in East Asia over the last deglaciation. *Earth Planet. Sci. Lett.* 301 (1–2), 256–264.
- Peterse, F., Martínez-García, A., Zhou, B., Beets, C.J., Prins, M.A., Zheng, H., Eglinton, T.I., 2014. Molecular records of continental air temperature and monsoon precipitation variability in East Asia spanning the past 130,000 years. *Quat. Sci. Rev.* 83, 76–82.
- Pirrung, M., Fischer, C., Buchel, G., Gaupp, R., Lutz, H., Neuffer, F.O., 2003. Lithofacies succession of maar crater deposits in the Eifel area (Germany). *Terra Nova* 15, 125–132.
- Ramrath, A., Zolitschka, B., Wulf, S., Negendank, J.F.W., 1999. Late Pleistocene climatic variations as recorded in two Italian maar lakes (Lago di Mezzano, Lago Grande di Monticchio). *Quat. Sci. Rev.* 18 (7), 977–992.
- Ramrath, A., Sadori, L., Negendank, J.F.W., 2000. Sediments from Lago di Mezzano, central Italy: a record of Lateglacial/Holocene climatic variations and anthropogenic impact. *The Holocene* 10 (1), 87–95.
- Rasmussen, S.O., Andersen, K.K., Svensson, A.M., Steffensen, J.P., Vinther, B.M., Clausen, H.B., Siggaard-Andersen, M.L., Johnsen, S.J., Larsen, L.B., Dahl-Jensen, D., Bigler, M., Röthlisberger, R., Fischer, H., Goto-Azuma, K., Hansson, M.E., Ruth, U., 2006. A new Greenland ice core chronology for the last glacial termination. *J. Geophys. Res.* 111 (D6).
- Rasmussen, S.O., Seierstad, I.K., Andersen, K.K., Bigler, M., Dahl-Jensen, D., Johnsen, S.J., 2008. Synchronization of the NGRIP, GRIP, and GISP2 ice cores across MIS 2 and palaeoclimatic implications. *Quat. Sci. Rev.* 27 (1–2), 18–28.
- Reichgelt, T., Kennedy, E.M., Mildenhall, D.C., Conran, J.G., Greenwood, D.R., Lee, D.E., 2013. Quantitative palaeoclimate estimates for Early Miocene southern New Zealand: evidence from Foulden Maar. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 378, 36–44.

- Richter, G., Baszio, S., Wuttke, M., 2017. Discontinuities in the microfossil record of middle Eocene Lake Messel: clues for ecological changes in lake's history? *Palaeobiodivers. Palaeoenviron.* 97 (2), 295–314.
- Rioual, P., Andrieu-Ponel, V., Beaulieu, J.-L., Reille, M., Svobodova, H., Battarbee, R.W., 2007. Diatom responses to limnological and climatic changes at Ribains Maar (French Massif Central) during the Eemian and Early Würm. *Quat. Sci. Rev.* 26, 1557–1609.
- Ruiz-Fernández, A.C., Hillaire-Marcel, C., Páez-Osuna, F., Ghaleb, B., Caballero, M., 2007. 210 Pb chronology and trace metal geochemistry at Los Tuxtlas, Mexico, as evidenced by a sedimentary record from the Lago Verde crater lake. *Quat. Res.* 67 (2), 181–192.
- Sabol, M., Vass, D., Konecny, V., Hudackova-Hlavata, N., Slamkova, M., Tunyi, I., Balogh, K., 2004. Reconstruction of the paleoenvironmental evolution of the Bone Gorge Maar near Hajnacka (southern Slovakia). Occasional Papers of the Geological Institute of Hungary. 203, pp. 88–89.
- Schettler, G., Liu, Q., Mingram, J., Negendank, J.F.W., 2006a. Palaeovariations in the East-Asian Monsoon regime geochemically recorded in varved sediments of Lake Sihailongwan (Northeast China, Jilin Province). Part 1: hydrological conditions and dust flux. *J. Paleolimnol.* 35 (2), 239–270.
- Schettler, G., Liu, Q., Mingram, J., Stebich, M., Dulski, P., 2006b. East-Asian monsoon variability between 15,000 and 2000 cal. yr BP recorded in varved sediments of Lake Sihailongwan (northeastern China, Long Gang volcanic field). *The Holocene* 16 (8), 1043–1057.
- Schettler, G., Mingram, J., Negendank, J.F.W., Jiaqi, L., 2006c. Palaeovariations in the east-Asian monsoon regime geochemically recorded in varved sediments of Lake Sihailongwan (Northeast China, Jilin Province). Part 2: a 200-year record of atmospheric Lead-210 flux variations and its palaeoclimatic implications. *J. Paleolimnol.* 35 (2), 271–288.
- Schulz, M., Berger, W.H., Sarnthein, M., Grootes, P.M., 1999. Amplitude variations of 1470-year climate oscillations during the last 100,000 years linked to fluctuations of continental ice mass. *Geophys. Res. Lett.* 26 (22), 3385–3388.
- Sikes, E.L., Medeiros, P.M., Augustinus, P., Wilmshurst, J.M., Freeman, K.R., 2013. Seasonal variations in aridity and temperature characterize changing climate during the last deglaciation in New Zealand. *Quat. Sci. Rev.* 74 (74), 245–256.
- Sirocko, F., 2016. The ELSA - stacks (Eifel-Laminated-Sediment-Archive): an introduction. *Glob. Planet. Chang.* 142, 96–99.
- Sirocko, F., Dietrich, S., Veres, D., Grootes, P.M., Schaber-Mohr, K., Seelos, K., Nadeau, M.J., Kromer, B., Rothacker, L., Röhrner, M., 2013. Multi-proxy dating of Holocene maar lakes and Pleistocene dry maar sediments in the Eifel, Germany. *Quat. Sci. Rev.* 62 (62), 56–76.
- Stebich, M., Brühmann, C., Kulbe, T., Negendank, J.F.W., 2005. Vegetation history, human impact and climate change during the last 7000 years recorded in annually laminated sediments of Lac Pavin, France. *Rev. Palaeobot. Palynol.* 133 (1), 115–133.
- Stebich, M., Mingram, J., Han, J.T., Liu, J.Q., 2009. Late Pleistocene spread of (cool-)temperate forests in Northeast China and climate changes synchronous with the North Atlantic region. *Glob. Planet. Chang.* 65 (1–2), 56–70.
- Stebich, M., Rehfeld, K., Schlütz, F., Tarasov, P.E., Liu, J., Mingram, J., 2015. Holocene vegetation and climate dynamics of NE China based on the pollen record from Sihailongwan Maar Lake. *Quat. Sci. Rev.* 124, 275–289.
- Sun, Q., Chu, G., Shengqiang, L.L., 2004. Long-chain alkenones in sulfate lakes and its paleoclimatic implications. *Chin. Sci. Bull.* 49 (19), 2082–2086 (in Chinese with English abstract).
- Sun, Q., Chu, G., Liu, G., Li, S., Wang, X., 2007. Calibration of alkenone unsaturation index with growth temperature for a lacustrine species, *Chrysotila lamellosa* (Haptophyceae). *Org. Geochem.* 38 (8), 1226–1234.
- Sun, Q., Chu, G., Liu, M., Xie, M., Li, S., Ling, Y., Wang, X., Shi, L., Jia, G., Lü, H., 2011. Distributions and temperature dependence of branched glycerol dialkyl glycerol tetraethers in recent lacustrine sediments from China and Nepal. *J. Geophys. Res.* 116 (G1).
- Sun, C., Plunkett, G., Liu, J., Zhao, H., Sigl, M., McConnell, J.R., Pilcher, J.R., Vinther, B., Steffensen, J.P., Hall, V., 2014. Ash from Changbaishan Millennium eruption recorded in Greenland ice: implications for determining the eruption's timing and impact. *Geophys. Res. Lett.* 41 (2), 694–701.
- Sun, C., You, H., He, H., Zhang, L., Gao, J., Guo, W., Chen, S., Mao, Q., Liu, Q., Chu, G., 2015. New evidence for the presence of Changbaishan Millennium eruption ash in the Longgang volcanic field, Northeast China. *Gondwana Res.* 28 (1), 52–60.
- Sun, C., Liu, J., You, H., Chu, G., 2016a. Clinopyroxene and Fe-Ti oxides for correlating the ash from Changbaishan Millennium eruption. *Sci. China Earth Sci.* 59 (7), 1454–1462.
- Sun, Q., Shan, Y., Sein, K., Su, Y., Zhu, Q., Wang, L., Sun, J., Gu, Z., Chu, G., 2016b. A 530 year long record of the Indian Summer Monsoon from carbonate varves in Maar Lake Twintaung, Myanmar. *J. Geophys. Res. Atmos.* 121 (10), 5620–5630.
- Sun, C., Wang, L., Plunkett, G., You, H., Zhu, Z., Zhang, L., Zhang, B., Chu, G., Liu, J., 2018. Ash From the Changbaishan Qixiangshan Eruption: A New Early Holocene Marker Horizon Across East Asia. *J. Geophys. Res. Solid Earth* <https://doi.org/10.1029/2018JB015983> in press.
- Tao, K.Y., 2007. Leiqiong Global Geopark in China. *Resour. Surv. Environ.* 28 (3), 223–227 (in Chinese with English abstract).
- Telford, R.J., Lamb, H.F., 1999. Groundwater-mediated response to holocene climatic change recorded by the diatom stratigraphy of an Ethiopian crater lake. *Quat. Res.* 52 (1), 63–75.
- Tierney, J.E., Russell, J.M., Eggermont, H., Hopmans, E.C., Verschuren, D., Sissingh Damsté, J.S., 2010. Environmental controls on branched tetraether lipid distributions in tropical East African lake sediments. *Geochim. Cosmochim. Acta* 74 (17), 4902–4918.
- Toney, J.L., Huang, Y., Fritz, S.C., Baker, P.A., Grimm, E., Nyren, P., 2010. Climatic and environmental controls on the occurrence and distributions of long chain alkenones in lakes of the interior United States. *Geochim. Cosmochim. Acta* 74 (5), 1563–1578.
- Turner, C., 1998. Volcanic maars, long Quaternary sequences and the work of the INQUA subcommission on European Quaternary stratigraphy. *Quat. Int.* 47–48, 41–49.
- Valentine, G.A., 2012. Shallow plumbing systems for small-volume basaltic volcanoes, 2: evidence from crustal xenoliths at scoria cones and maars. *J. Volcanol. Geotherm. Res.* 223–224, 47–63.
- Valentine, G.A., White, J.D.L., Ross, P.S., Graetinger, A.H., Sonder, I., 2017. Updates to concepts on phreatomagmatic maar-diatremes and their pyroclastic deposits. *Front. Earth Sci.* 5, 68.
- Vass, D., Konecny, V., Tunyi, I., Dolinsky, P., Balogh, K., Hudackova, N., Kovacova-Slamkova, M., Belacek, B., 2000. Origin of the Pliocene vertebrate bone accumulation at Hajnacka, southern Slovakia. *Geol. Carpath.* 51 (2), 69–82 (Bratislava).
- Vos, H., Sanchez, A., Zolitschka, B., Brauer, A., Negendank, J.F.W., 1997. Solar activity variations recorded in varved sediments from the crater lake of Holzmaar – a maar lake in the Westeifel Volcanic Field, Germany. *Surv. Geophys.* 18 (2–3), 163–182.
- Vos, H., Mingram, J., Negendank, J.F.W., 2000. Comparative time series analyses of laminated sediments from central European maar lakes (Eckfeld + Messel, Eocene; Pula + Gerce, Pliocene). *Terra Nostra* 6, 485–496.
- Walker, D., 2007. Holocene sediments of Lake Barrine, north-east Australia, and their implications for the history of lake and catchment environments. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 251 (1), 57–82.
- Walker, D., 2011. The frequency of lamination in the sediments of Lake Barrine, tropical north-east Australia, during the last five millennia. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 299 (1), 214–226.
- Walker, D., Owen, J.A.K., 1999. The characteristics and source of laminated mud at Lake Barrine, Northeast Australia. *Quat. Sci. Rev.* 18 (14), 1597–1624.
- Wang, P., Tian, J., Cheng, X., Liu, C., Xu, J., 2003. Carbon reservoir changes preceded major ice-sheet expansion at the mid-Brunhes event. *Geology* 31 (3), 239–242.
- Wang, S.Y., Lü, H.Y., Liu, J.Q., Negendank, J.F.W., 2007. The early Holocene optimum inferred from a high-resolution pollen record of Huguangyan Maar Lake in southern China. *Chin. Sci. Bull.* 52 (20), 2829–2836.
- Wang, L., Lu, H., Liu, J., Gu, Z., Mingram, J., Chu, G., Li, J., Rioual, P., Negendank, J.F.W., Han, J., 2008. Diatom-based inference of variations in the strength of Asian winter monsoon winds between 17,500 and 6,000 calendar years B.P. *J. Geophys. Res. Atmos.* 113 (D21), 6089–6098.
- Wang, L., Li, J., Lu, H., Gu, Z., Rioual, P., Hao, Q., Mackay, A.W., Jiang, W., Cai, B., Xu, B., Han, J., Chu, G., 2012. The East Asian winter monsoon over the last 15,000 years: its links to high-latitudes and tropical climate systems and complex correlation to the summer monsoon. *Quat. Sci. Rev.* 32, 131–142.
- Wang, X., Chu, G., Sheng, M., Zhang, S., Li, J., Chen, Y., Tang, L., Su, Y., Pei, J., Yang, Z., 2016. Millennial-scale Asian summer monsoon variations in South China since the last deglaciation. *Earth Planet. Sci. Lett.* 451, 22–30.
- Wang, M., Zheng, Z., Man, M., Hu, J., Gao, Q., 2017. Branched GDGT-based paleotemperature reconstruction of the last 30,000 years in humid monsoon region of Southeast China. *Chem. Geol.* 463, 94–102.
- Wappler, T., Labandeira, C.C., Rust, J., Frankenhaeuser, H., Wilde, V., 2011. Testing for the effects and consequences of mid-Paleogene climate change on insect herbivory. Conference Volume - International Senckenberg Conference 22, pp. 168–169.
- Wastegård, S., Veres, D., Kliem, P., Hahn, A., Ohlendorf, C., Zolitschka, B., 2013. Towards a late Quaternary tephrochronological framework for the southernmost part of South America—the Laguna Potrok Aike tephra record. *Quat. Sci. Rev.* 71, 81–90.
- Weijers, J.W.H., Schefuss, E., Schouten, S., Damsté, J.S.S., 2007a. Coupled thermal and hydrological evolution of tropical Africa over the last deglaciation. *Science* 315 (5819), 1701–1704.
- Weijers, J.W.H., Schouten, S., van den Donker, J.C., Hopmans, E.C., Sissingh Damsté, J.S., 2007b. Environmental controls on bacterial tetraether membrane lipid distribution in soils. *Geochim. Cosmochim. Acta* 71 (3), 703–713.
- White, J., Houghton, B., 2006. Primary volcanoclastic rocks. *Geology* 34 (8), 677–680.
- White, J.D.L., Ross, P.S., 2011. Maar-diatreme volcanoes: a review. *J. Volcanol. Geotherm. Res.* 201, 1: 1–29.
- Wilkins, D., De Deckker, P., Fifield, L.K., Gouramanis, C., Olley, J., 2012. Comparative optical and radiocarbon dating of laminated Holocene sediments in two maar lakes: Lake Keilambete and Lake Gnotuk, south-western Victoria, Australia. *Quat. Geochronol.* 9, 3–15.
- Williams, T., Thouveny, N., Creer, K., 1998. A normalised intensity record from Lac du Bouchet: geomagnetic palaeointensity for the last 300 kyr? *Earth Planet. Sci. Lett.* 156 (1–2), 33–46.
- Williamson, D., Jelinowska, A., Kissel, C., Tucholka, P., Gibert, E., Gasse, F., Massault, M., Taieb, M., Van Campo, E., Wieckowski, K., 1998. Mineral-magnetic proxies of erosion/oxidation cycles in tropical maar-lake sediments (Lake Tritrivalakely, Madagascar): paleoenvironmental implications. *Earth Planet. Sci. Lett.* 155 (3), 205–219.
- Willis, K.J., Braun, M., 1997. Terrestrial responses to Milankovitch Cyclicity in the late Pliocene; evidence from the laminated sediments of the Pula Maar Crater Lake, Hungary. *Wuerzburger Geographische Manuskripte.* 41, p. 229.
- Willis, K.J., Kleczkowski, A., Briggs, K.M., Gilligan, C.A., 1999. The role of sub-Milankovitch climatic forcing in the initiation of the Northern Hemisphere glaciation. *Science* 285 (5427), 568–571.
- Wulfs, S., Kraml, M., Brauer, A., Keller, J., Negendank, J.F., 2004. Tephrochronology of the 100 ka lacustrine sediment record of Lago Grande di Monticchio (southern Italy). *Quat. Int.* 122 (1), 7–30.
- Yamada, K., Kamite, M., Saito-Kato, M., Okuno, M., Shinozuka, Y., Yasuda, Y., 2010. Late Holocene monsoonal-climate change inferred from Lakes Ni-no-Megata and Sano-Megata, northeastern Japan. *Quat. Int.* 220 (1), 122–132.

- Yancheva, G., Nowaczyk, N.R., Mingram, J., Dulski, P., Schettler, G., Negendank, J.F.W., Liu, J., Sigman, D.M., Peterson, L.C., Haug, G.H., 2007. Influence of the intertropical convergence zone on the East Asian monsoon. *Nature* 445 (7123), 74–77.
- Yang, X., Heller, F., Yang, J., Su, Z., 2009. Paleosecular variations since ~9000 yr BP as recorded by sediments from maar lake Shuangchiling, Hainan, South China. *Earth Planet. Sci. Lett.* 288 (1–2), 1–9.
- You, H.T., Liu, J.Q., 2012. High-resolution climate evolution derived from the sediment records of Erlongwan Maar Lake since 14 ka BP. *Sci. Bull.* 57 (27), 3610–3616.
- You, H., Liu, J., Liu, Q., Chu, G., Rioual, P., Han, J., 2008. Study of the varve record from Erlongwan maar lake, NE China, over the last 13 ka BP. *Chin. Sci. Bull.* 53 (2), 262–266.
- Zaarur, S., Stein, M., Adam, O., Mingram, J., Chu, G., Liu, J., Wu, J., Erel, Y., 2018. Late Quaternary climate in southern China deduced from Sr–Nd isotopes of Huguangyan Maar sediments. *Earth Planet. Sci. Lett.* 496, 10–19.
- Zheng, Z., Guiot, J., 1999. A 400,000-year paleoclimate reconstruction in tropical region of China. *Acta Sci. Natur. Univ. Sunyatseni* 38 (6), 94–98 (in Chinese with English abstract).
- Zheng, Z., Lei, Z.Q., 1999. A 400,000 year record of vegetational and climatic changes from a volcanic basin, Leizhou Peninsula, southern China. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 145 (4), 339–362.
- Zheng, Y., Pancost, R.D., Liu, X., Wang, Z., Naafs, B.D.A., Xie, X., Liu, Z., Yu, X., Yang, H., 2017. Atmospheric connections with the North Atlantic enhanced the deglacial warming in northeast China. *Geology* 45 (11), 1031–1034.
- Zink, K.-G., Vandergoes, M.J., Mangelsdorf, K., Dieffenbacher-Krall, A.C., Schwark, L., 2010. Application of bacterial glycerol dialkyl glycerol tetraethers (GDGTs) to develop modern and past temperature estimates from New Zealand lakes. *Org. Geochem.* 41 (9), 1060–1066.
- Zink, K.G., Vandergoes, M.J., Bauersachs, T., Newnham, R.M., Rees, A.B.H., Schwark, L., 2016. A refined paleotemperature calibration for New Zealand limnic environments using differentiation of branched glycerol dialkyl glycerol tetraether (brGDGT) sources. *J. Quat. Sci.* 31 (7), 823–835.
- Zolitschka, B., 1998. A 14,000 year sediment yield record from western Germany based on annually laminated sediments. *Geomorphology* 22 (1), 1–17.
- Zolitschka, B., Negendank, J.F.W., Lottermoser, B.G., 1995. Sedimentological proof and dating of the Early Holocene volcanic eruption of Ulmener Maar (Vulkaneifel, Germany). *Geol. Rundsch.* 84 (1), 213–219.
- Zolitschka, B., Brauer, A., Negendank Jr., F.W., Stockhausen, H., Lang, A., 2000. Annually dated late Weichselian continental paleoclimate record from the Eifel, Germany. *Geology* 28 (9), 783–786.
- Zolitschka, B., Anselmetti, F., Ariztegui, D., Corbella, H., 2009. The Laguna Potrok Aike scientific drilling project PASADO (ICDP Expedition 5022). *Sci. Drill.* 8 (8), 75–82.
- Zolitschka, B., Anselmetti, F., Ariztegui, D., Corbella, H., Francus, P., Lücke, A., Maidana, N.I., Ohlendorf, C., Schäbitz, F., Wastegård, S., 2013. Environment and climate of the last 51,000 years – new insights from the Potrok Aike maar lake Sediment Archive Drilling project (PASADO). *Quat. Sci. Rev.* 71, 1–12.
- Zolitschka, B., Francus, P., Ojala, A.E., Schimmelmänn, A., 2015. Varves in lake sediments—a review. *Quat. Sci. Rev.* 117, 1–41.